



US007867721B2

(12) **United States Patent**  
**Zuker et al.**

(10) **Patent No.:** **US 7,867,721 B2**  
(45) **Date of Patent:** **\*Jan. 11, 2011**

(54) **IN VITRO METHOD FOR IDENTIFYING A COMPOUND USING AN EUKARYOTIC MECHANOSENSORY TRANSDUCTION CHANNEL**

(75) Inventors: **Charles S. Zuker**, San Diego, CA (US);  
**Richard G. Walker**, La Jolla, CA (US);  
**Aarron Willingham**, La Jolla, CA (US)

(73) Assignee: **The Regents of the University of California**, Oakland, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 552 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **11/703,343**

(22) Filed: **Feb. 6, 2007**

(65) **Prior Publication Data**

US 2009/0075294 A1 Mar. 19, 2009

**Related U.S. Application Data**

(60) Continuation of application No. 10/369,978, filed on Feb. 18, 2003, now abandoned, which is a division of application No. 09/392,812, filed on Sep. 9, 1999, now Pat. No. 6,537,778.

(51) **Int. Cl.**

**G01N 33/53** (2006.01)  
**C12N 5/00** (2006.01)  
**C07K 14/00** (2006.01)

(52) **U.S. Cl.** ..... **435/7.2; 435/325; 530/350**

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,824,306 A 10/1998 Tang et al.

**OTHER PUBLICATIONS**

Corey, et al., New TRP channels in hearing and mechanosensation, 2003, Cell, vol. 39, pages 585-588.

Garcia-Anoveros, J, et al. "The Molecules of Mechanosensation," Annu. Rev. Neurosci., 1997, 20:567-594.

Gillespie, et al., "Have we found the tip link, transduction channel, and gating spring of the hair cell?" 2005, Current Opinion in Neurobiology, vol. 15, pp. 389-396.

Hudspeth, A.J., "How the ear's works work," nature, 1989, 341:397-404.

Kernan, M. et al., "Genetic Dissection of Mechanosensory Transduction: Mechanoreception-Defective Mutations of Drosophila," Neuron, 1994, 12:1195-1206.

Liman, et al. "TRP2: A candidate transduction channel for mammalian pheromone sensory signaling." Proc. Natl. Acad. Sci., USA. May 1999, vol. 96, pp. 5791-5796.

Mikayama, T. "Molecular cloning and functional expression of a cDNA encoding glycosylation-inhibiting factor." Proc. Natl. Acad. Sci. USA vol. 90, pp. 10056-10060, 1993.\*

Montell, C., "TRP trapped in fly signaling web," Current Opinion in Neurobiology, 8:389-397 (1998).

Sukharev, S.I., et al., "A large-conductance mechanosensitive channel in *E. coli* encoded by *mscL* alone," nature, 1994, 368:265-268.

Vannier, et al. Mouse *trp2*, the homologue of the human *trpc2* pseudogene, encodes *mTrp2*, a store depletion-activated capacitative Ca<sup>2+</sup> entry channel. Proc. Natl. Acad. Sci. USA. Mar. 1999, vol. 96, pp. 2060-2064.

Voet, et al. Biochemistry. 1990. John Wiley & Sons, Inc.. pp. 126-128 and 228-234.

Wissenbach, et al. Structure and mRNA expression of a bovine *trp* homologue related to mammalian *trp2* transcripts. FEBS letters. 1998, vol. 429, pp. 61-66.

*Primary Examiner*—Robert Landsman

*Assistant Examiner*—Ian Dang

(74) *Attorney, Agent, or Firm*—Townsend and Townsend and Crew LLP

(57) **ABSTRACT**

The present invention provides, for the first time, nucleic acids encoding a eukaryotic mechanosensory transduction channel (MSC) protein. The proteins encoded by these nucleic acids form channels that can directly detect mechanical stimuli and convert them into electrical signals. These nucleic acids and the proteins they encode can be used as probes for sensory cells in animals, and can be used to diagnose and treat any of a number of human conditions involving inherited, casual, or environmentally-induced loss of mechanosensory transduction activity.

**7 Claims, 3 Drawing Sheets**

FIG. 1A

```

Drosophila -----
C.elegans MSRSEKCLTVRKRET RSTSVTRAEMFTGKK MDAAKNAFDLLTTDT IPTDRPPLRRSSTHL QIGKNSRIIFVPKQP SRDSVTPPDRLLGKP -----

Drosophila -----
C.elegans LFRESLTSHASSHEE MSSEDLAMADPQTKI LYFAKRDEWANVESE IETIKRSDFSMADNH GFTAFLLAVKAGKDQ IVDKMIRKGARVDYS -----

Drosophila -----
C.elegans TKOGRNATHIAAMYS GVETLELILKRYSEL LRGAGPKKQLAIHV ACERKSKKAFPIVKR ILEDTDQRMAEDGDG SLPIHLAFKFGWVNI -----

Drosophila -----
C.elegans VELLSSGSPDEQTRK ADNGDTHLLHLAARS GNIEAVRTAIAAGCD NANVQNRVGRTPLHE CLTVTGTQKGYVAEV GDQNMLKIMFKLRAD -----

Drosophila -----
C.elegans -----RTPMHLA AENGHAHVIEILADK FKASIFERTKDGSTL MHIASLNHGAECATM LFKKGVYVHMPNKDG ARSIHTAAAYGHTGI -----
ANIHKEDKTPVHVA AERGDTSMWESLIDK FGGIRARTRDGSSTL LHIAACSGHTSTALA FLKR-VPLFMPNKKG ALGLHSAAAAAGFNDV -----

Drosophila -----
C.elegans INTLLQKGEKVDVTT -NNYTALHIAVESAK PAVVETLLGFGADVH VRGGKLRRETPLHIAA RVKGDRCALMLLKS GASPNTTDDCLTPV -----
VKMLIARGTNDVVRT RDNYTALHVAVQSGK ASVVETLLGSGADIH VKGG---E----- -LMDGETC-----L -----

Drosophila -----
C.elegans HVAARHGNLATLMQL LEDEGPLYKSNTGE TPLHMACRACHPDIV RHLIETVKEKHGPKD ATTYNSVNEEDGATA LHYTCQITKEEVKIP -----
HIAARSGNKD-IMLL LDENADSKISSKIGE TPLQVAAKSCNFEEA SMILKHLSEVLTQEQ LKEHVNHRTNDGFTA LHYYAAEIEQRQLHFP -----

Drosophila -----
C.elegans ESDKQIVRMILLENGA DVTLQTKTALETAFH YCAVAGNNDVLEMI SHMNPTDIQKAMNRQ SSVGWTPLLIACHRQ HMELVNNLL-----
GEDAKLVNLLIDYGG MVEMPSLNANETAMH MAARSGNQAVLLAMV NKIGAGAVQIVQNKQ SKNGWSPLLEACARG HSGVANILLKVLVLC -----

Drosophila -----
C.elegans VGPGPGPRLQGRG YWTRTRARVTPWLQ YQGYWARTRIRTRAR ATGPGLDQDQGYWART RTRTKVTPRLLGDH HARIDVDFDEMGRAL -----
-----AN HARVDVDFTEGRSAL -----

Drosophila -----
C.elegans HLAALERGYLHVCDAL LTNKAFINSKSRVGR TALHLAAMNGFTHLV KFLIKDHNVIDILT LRKQTPHLAAASGQ MEVCQLLLELGANID -----
HLAAFNGHLSLVHLL LQHKAFVNSKSKTGE APLHLAAQHGHVKYV NVLVQDGHGALEAIT LDNQ TALHFAAKFGQ LAVSQTLALGANPN -----
    
```

FIG. 1B

<i>Drosophila</i>	ATDDLQKPIHVAQ	NNYSEVAKLFLQQHP	---SLVNATSKDNT	CAHIAAMQGSVKVTE	ELMKFDRSGVISARN	KLTDATPLQLAAEGG
<i>C.elegans</i>	ARDDKGQTPHLAAE	NDFPDVVKLFLKMRN	NRSVLTAIDHNGFT	CAHIAAMKGS LAVVR	ELMMIDKPMVIQAKT	KTLEATTLLHMAAAGG
<i>Drosophila</i>	HADVVKALVRAGASC	TEENKAGFTAVHLAA	QNGHGQVLDVLKSTN	SLRINSKKGGLTPLH	VAAYYQADTVRELL	TSPATVKSETPTG--
<i>C.elegans</i>	HANIVKILLENGANA	ELEN-SGMTALHLGA	KNGFISILEAFDKIL	WKRC-SRKTGLNALH	IAAFYGNDFVNEML	KHYQATVRSEPPYYN
<i>Drosophila</i>	QSLFGDLGTESGMTP	LHLAAFSGNENVVR	LLNSAGVQVDAATIE	NMHGHIQMV EILLGQ	GAEINATDRNGWTP	HCAAKAGHLEVVKLL
<i>C.elegans</i>	HVNKEFSTEYGF	LHLAAHSGHDSLVRM	LLN-QGVQVDATSTT	MMS-----	-----	-----
<i>Drosophila</i>	CEAGASPKSETNYGC	AAIMFAASEGHNEVL	RYLMNKEHDTYGLME	DKRFVYNLMVSKNH	NNKPIQEFVLVSPAP	VDTAAKLSNIYIVLS
<i>C.elegans</i>	-----	-----	-----	-----K-----	-----E-----	-----E-----
<i>Drosophila</i>	TKKERAKDLVAAGKQ	CEAMATELLALAAGS	DSAGKILQATDKRNV	EFLDVLIENEQKEVI	AHTVVQRYLQELWHG	SLTWASWKILLLLVA
<i>C.elegans</i>	--KERAKDLLNVAVF	SEMMAVELLITATEY	N-AALLLKAKDNRGR	PLLDVLIENEQKEVW	SYASVQRYL TEVWTA	RVDWSFGKFVAFSLF
<i>Drosophila</i>	FIVCPPVWIGFTFPM	GHKFNKVP IIKFMSY	LTSHIYMIHLSIVG	ITPIYVLRLSLVPY	WYEVGLLIWL SGLLL	FELTNPSDKSGLGSI
<i>C.elegans</i>	VLICPPAWFYFSLPL	DSRIGRAPIIKFVCH	IVSHVYFTILLTIIV	LNITHKYEVT SVVPN	PVEMLLLLWLSGNLV	SELSTVGGGSGLGIV
<i>Drosophila</i>	KVLVLLGMAGVGVH	VSAFLV S-----	-----KEYWPTLVYCR	NQCFALAFLLACVQI	LDFLSFHHLFGPWAI	IIGDLLKDLARFLAV
<i>C.elegans</i>	KVLILVLSAMAIAVH	VLAFLLPVFLTHLD	NDEKLFARTMLYLK	NQLFAFALLFAFVEY	LDFLT VHHLFGPWAI	II---MYDLARFLVI
<i>Drosophila</i>	LAI FVFGFSMHIVAL	NQSFANFSPEDLRSF	EKKNRNRGYFSDMEQ	MTCPHDLRRRIMS	IVASANSDESTRTTF	PGGTSTSPHSLL EIP
<i>C.elegans</i>	LMLFVAGFTLHVTSI	-----FQP-----	-----	-----AYQPVE-----	-----	-----
<i>Drosophila</i>	SPCMHVDVFIQSIQT	KIKQISINIDITNAR	LPGAFSLRRLPTTKF	CTIETIETDRIESIT	KDNATD TDYRCSYM	LGPMT PFLAFERLFF
<i>C.elegans</i>	-----	-----	-----	---DSAELMRLASPS Q-----	-----	-----TLEMLFF
<i>Drosophila</i>	AVFGQTTTLDINPMR	HLRPEVTEVLFKFFV	GIYLLVSVVVLINLL	IAMMSDTYQRIQMN-	--RNHGLVDRTNQRNK	KKKKNHIIESTNPTW
<i>C.elegans</i>	SLFGLVEP-DSMPPL	HLVPDFAKIILKLLF	GIYMMVTLIVLINLL	IAMMSDTYQRIQAQS	DKENKFGRAILIRQM	NKK-----SATPSP

FIG. 1C

Drosophila	ASVIFLFFKIISTPA	NICVLSGGVYLYLYL	YLEMYLWVSDTVRMH	PINSFELLFFAVFGQ	TTTEQTQVDKIKNVA	TPTQPYWVEYLFKIV
C.elegans	INMLTKLIIVLRVAW	RNRGK-APLSTPLAS	FRCMTRKAQDDLRFE	EN---IDAFSMGGGQ	-----QGRQ	SPTNEGRGQ-----
Drosophila	FGIYMLVSVVVLINL	LIAMMSDYYQRIQAQ	SDIEWKFGLSKLI RN	MHRTTTAPSPNLVT	TWFMWIVEKVKVKSQ	VTKVAFQPLSCLSL
C.elegans	-----QE--L	GNSADWN-----IE	TVIDWRKIVSMYYQA	NGKLTGRTKEDVD-	--LAMAVPTSF	IKPQ G-----PDTTCR--
Drosophila	SIRILYPVSYTCFHI	CMKKKRPSLVQMMG	IRQASPRTKAGAKWL	SKIKKSVALSQVHLS	PLGSQASFSQANQNR	IENVADWEAIAKKYR
C.elegans	-----PIDYTWLRL	CKTKS-----HG	SGLSIVRRKTRGKIV	YSTRTNTSVLQINSS	-----R-NAPK	IYLRYGRAKIAHFFF
Drosophila	ALVGDEEGGSLKDS	AESGSEQSGGQPP	AQVGRRAIKATLADT	TKSKLHLSLQITILPD	YLYLFSTIQASVLLC	TLGMVFSDSGTHFFW
C.elegans	TST-----TLKG--	---GAFMWHG-----	-----LAARLCKIRVD	HM-----	-----	-----
Drosophila	FNWSMGKSD	1704				
C.elegans	-----	1709				

**IN VITRO METHOD FOR IDENTIFYING A  
COMPOUND USING AN EUKARYOTIC  
MECHANOSENSORY TRANSDUCTION  
CHANNEL**

CROSS-REFERENCES TO RELATED  
APPLICATIONS

This application is a continuation of U.S. application Ser. No. 10/369,978, filed Feb. 18, 2003, now abandoned which is a divisional of U.S. application Ser. No. 09/392,812, filed Sep. 9, 1999, which is now U.S. Pat. No. 6,537,778. The contents of each are incorporated herein by reference in their entireties.

STATEMENT AS TO RIGHTS TO INVENTIONS  
MADE UNDER FEDERALLY SPONSORED  
RESEARCH AND DEVELOPMENT

This invention was made with Government support under Grant No. DC03160, awarded by the National Institutes of Health. The Government has certain rights in this invention.

FIELD OF THE INVENTION

This invention provides isolated nucleic acid and amino acid sequences of a novel family of eukaryotic mechanosensory ion channels that are designated mechanosensory transduction channels (MSC).

BACKGROUND OF THE INVENTION

The ability to detect mechanical stimuli is an essential and prevalent characteristic of living organisms, and is found from bacteria to simple metazoans to the most complex of mammals. Indeed, the ability to detect mechanical stimuli and convert them into electrical signals forms the basis of many central aspects of animal life, such as light touch, heavy touch, proprioception, baroreception, balance, and the crown jewel, hearing. Even the ability of cells to stop growing when in contact with neighboring cells is likely dependent on mechanical stimuli. Not surprisingly, therefore, numerous human conditions result at least in part from an inability to detect mechanical stimuli, such as Meniere's Disease, sensorineural deafness, blood pressure disorders, and various types of cancers.

In general, the variety of known mechanosensory modalities are thought to be mediated by mechanically-gated cation channels present within the membrane of receptor cells. This view has come in large part from detailed studies into the physiology of mechanosensation using various cell types involved in mechanosensory detection, such as the hair cells of the vertebrate inner ear, single-celled ciliates such as *Paramecium*, or the sensory neurons of *Drosophila* (see, e.g., Kernan et al., *Neuron* 12:1195-1206 (1994)). In *Drosophila*, the dendrite of the sensory neuron is enclosed in a cavity filled with a specialized receptor lymph, which is unusually rich in potassium ions, and is functionally equivalent to the potassium-rich endolymph of the vertebrate cochlea. These potassium ions produce a transepithelial potential difference, with the apical side of the epithelium being positively charged. Mechanical stimulation of the bristle, which is adjacent to the sensory neuron, generates a mechanoreceptor potential within the neuron, detectable as a negative deflection of the transepithelial potential, which reflects the flow of cations from the receptor lymph into the sensory neuron.

Activation of the hair cells of vertebrates also result in the influx of cations into cells (see, e.g., Hudspeth, *Nature*, 341: 397-404 (1989)). Each hair cell has a number of specialized microvillar structures, called stereocilia, whose deflection results in the activation of a putative channel present on the surface of the cell. Interestingly, electrophysiological studies have suggested that these cells contain a similar number of receptor channels as they do stereocilia, suggesting that perhaps each receptor channel is coupled to a single stereocilium. In addition, studies of the kinetics of hair-cell activation have suggested that the putative mechanosensory receptors are directly stimulated by mechanical force, resulting in the direct opening of the channel without the involvement of second messengers.

Despite the great importance of mechanosensation for animal behavior and health, and the detailed electrophysiological understanding that has been gained from the above-described studies, almost nothing is known about the molecular basis of mechanosensory detection in eukaryotes. Several mutations and distantly related molecules involved in this process have, however, been found. In *Drosophila*, for example, a number of mutations have been isolated that disrupt mechanoreception, resulting in a variety of phenotypes such as reduced locomotor activity, total uncoordination, and even death (Kernan et al., *Neuron* 12:1195-1206 (1994)). Also, mutations have been identified in the nematode *C. elegans* that result in a loss of sensitivity to gentle touch (reviewed in Garcia-Aanoveros & Corey, *Ann. Rev. Neurosci.* 20:567-594 (1997)). In addition, a prokaryotic mechanosensory channel has been identified (Sukarev et al., *Nature* 368: 265-268 (1994)). Still, despite these advances, the principle molecule of the mechanosensory transduction process in eukaryotes, the mechanically gated channel, has yet to be isolated or identified.

The identification and isolation of eukaryotic mechanosensory transduction channels would allow for the development of new methods of pharmacological and genetic modulation of mechanosensory transduction pathways. For example, availability of mechanosensory transduction channel proteins would permit screening for high-affinity agonists, antagonists, and modulators of mechanosensation in animals. Such molecules could then be used, e.g., in the pharmaceutical industry, to treat one or more of the many human conditions involving loss or hyperactivation of mechanosensation. In addition, the determination of nucleotide and amino acid sequences of mechanosensory transduction channels associated with a human condition would provide new tools for the diagnosis and/or treatment, e.g., gene-based treatment, of the condition.

SUMMARY OF THE INVENTION

The present invention provides for the first time nucleic acids encoding a eukaryotic mechanosensory transduction protein. The nucleic acids and the polypeptides they encode are referred herein as mechanosensory channel (MSC) nucleic acids and proteins. In vivo, MSC proteins form mechanosensory transduction channels that play a central role in many critical processes such as hearing, proprioception, and tactile sensation.

In one aspect, the present invention provides an isolated nucleic acid encoding a mechanosensory transduction protein, the protein having at least one of the following characteristics: (i) comprising greater than about 70% amino acid sequence identity to SEQ ID NO:2 or SEQ ID NO:4; (ii) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID

NO:9; or (iii) specifically binding to polyclonal antibodies generated against a polypeptide comprising an amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4; wherein the protein does not comprise the polypeptide sequence of SEQ ID NO:6.

In one embodiment, the nucleic acid encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4. In another embodiment, the nucleic acid comprises a nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3, but not SEQ ID NO:5.

In another embodiment, the nucleic acid selectively hybridizes under moderately stringent wash conditions to a nucleic acid comprising a nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3. In another embodiment, the nucleic acid selectively hybridizes under stringent wash conditions to a nucleic acid comprising a nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3, but not SEQ ID NO:5.

In another embodiment, the nucleic acid is amplified by primers that selectively hybridize under stringent hybridization conditions to the same sequence as degenerate primer sets encoding an amino acid sequence selected from the group consisting of: LDVLIENEQKEV (SEQ ID NO:7), HHLFGPWAI (SEQ ID NO:8), and VLINLLIAMMSD-TYQRIQ (SEQ ID NO:9).

In another embodiment, the nucleic acid is less than 120 kb. In another embodiment, the nucleic acid is less than 90 kb. In another embodiment, the nucleic acid is less than 60 kb. In another embodiment, the nucleic acid is less than 30 kb. In another embodiment, the nucleic acid is less than 10 kb. In another embodiment, the nucleic acid sequence encoding the MSC protein is isolated away from its genomic neighbors.

In another aspect, the present invention provides an expression cassette comprising a nucleic acid encoding a mechanosensory transduction protein, the protein having at least one of the following characteristics: (i) comprising greater than about 70% amino acid sequence identity to SEQ ID NO:2 or SEQ ID NO:4; (ii) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9; or (iii) specifically binding to polyclonal antibodies generated against a polypeptide comprising an amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4; wherein the protein does not comprise the polypeptide sequence of SEQ ID NO:6.

In another aspect, the present invention provides an isolated eukaryotic cell comprising the expression cassette.

In one aspect, the present invention provides an isolated nucleic acid encoding an extracellular domain of a mechanosensory transduction protein, the extracellular domain comprising greater than about 70% amino acid sequence identity to an extracellular domain of SEQ ID NO:2 or SEQ ID NO:4, wherein the extracellular domain does not comprise an extracellular domain of SEQ ID NO:6.

In one embodiment, the extracellular domain is fused to a heterologous polypeptide, thereby forming a chimeric polypeptide. In another embodiment, the extracellular domain comprises an amino acid sequence of an extracellular domain of SEQ ID NO:2 or SEQ ID NO:4.

In another aspect, the present invention provides an isolated mechanosensory transduction protein, the protein having at least one of the following characteristics: (i) comprising greater than about 70% amino acid sequence identity to SEQ ID NO:2 or SEQ ID NO:4; (ii) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9; or (iii) specifically binding to polyclonal antibodies generated against a polypeptide comprising an amino acid sequence of SEQ ID NO:2, or

SEQ ID NO:4; wherein the protein does not comprise the amino acid sequence of SEQ ID NO:6.

In one embodiment, the protein comprises the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4.

In another aspect, the present invention provides an isolated polypeptide comprising an extracellular domain of a mechanosensory transduction protein, the extracellular domain comprising greater than about 70% amino acid sequence identity to an extracellular domain of SEQ ID NO:2 or SEQ ID NO:4, wherein the extracellular domain does not comprise the amino acid sequence of an extracellular domain of SEQ ID NO:6.

In one embodiment, the extracellular domain is fused to a heterologous polypeptide, forming a chimeric polypeptide.

In another embodiment, the extracellular domain comprises the amino acid sequence of an extracellular domain of SEQ ID NO:2 or SEQ ID NO:4.

In another aspect, the present invention provides an antibody that selectively binds to a mechanosensory transduction protein, the protein having at least one of the following characteristics: (i) comprising greater than about 70% amino acid sequence identity to SEQ ID NO:2 or SEQ ID NO:4; (ii) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9; or (iii) specifically binding to polyclonal antibodies generated against a polypeptide comprising an amino acid sequence of SEQ ID NO:2, or SEQ ID NO:4; wherein the protein does not comprise the amino acid sequence of SEQ ID NO:6.

In another aspect, the present invention provides a method for identifying a compound that modulates mechanosensory receptor activity in eukaryotic cells, the method comprising the steps of: (i) contacting the compound with a mechanosensory receptor protein, the protein having at least one of the following characteristics: (a) comprising greater than about 70% amino acid sequence identity to a sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, and SEQ ID NO:6; (b) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9; or (c) specifically binding to polyclonal antibodies generated against a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, and SEQ ID NO:6; and (ii) determining the functional effect of the compound on the mechanosensory receptor protein.

In one embodiment, the mechanosensory receptor protein is expressed in a eukaryotic cell or cell membrane. In another embodiment, the functional effect is determined by detecting a change in the mechanoreceptor potential of the cell or cell membrane. In another embodiment, the functional effect is determined by detecting a change in an intracellular ion concentration. In another embodiment, the ion is selected from the group consisting of  $K^+$  and  $Ca^{2+}$ . In another embodiment, the protein comprises an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, and SEQ ID NO:6. In another embodiment, the protein is recombinant. In another embodiment, the functional effect is a physical interaction with the receptor protein.

In another aspect, the present invention provides a method of genotyping a human for a mechanosensory transduction channel locus, the method comprising detecting a mutation in a nucleic acid encoding a mechanosensory transduction channel in the human, the protein having at least one of the following characteristics: (a) comprising greater than about 70% amino acid sequence identity to a polypeptide having a sequence of SEQ ID NO:2; (b) having greater than about 90% amino acid sequence identity to a polypeptide having a

sequence of SEQ ID NO:5; (c) comprising an amino acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8; or (d) specifically binding to polyclonal antibodies generated against a polypeptide selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8; wherein the mutation introduces a premature stop codon into the nucleic acid 5' to the transmembrane domain region of the protein, or is a missense mutation removing a cysteine residue between transmembrane segments 4 and 5 of the protein.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A-C shows a continuing alignment between *Drosophila melanogaster* and *Caenorhabditis elegans* MSC homologs.

#### DETAILED DESCRIPTION OF THE INVENTION

##### Introduction

The present invention provides, for the first time, nucleic acids encoding a eukaryotic mechanosensory transduction channel (MSC) protein. Mutations in these nucleic acids and the proteins they encode are responsible for the "no-mechanoreceptor potential" phenotype in *Drosophila*, a phenotype involving uncoordination, often to the point of lethality, and a loss of mechanoreceptor potential in the bristles of mutant flies (Kernan et al., *Neuron* 12:1195-1206 (1994)). The proteins encoded by these nucleic acids form channels (e.g., as tetramers) that can directly detect mechanical stimuli and convert them into electrical signals. These proteins can detect mechanical stimuli in any of a number of sensory cells, such as neuronal sensory cells, hair cells, and others. These nucleic acids and the proteins they encode can be used as probes for sensory cells in animals, and can be used to diagnose and treat any of a number of human conditions involving inherited, casual, or environmentally-induced loss of mechanosensory transduction activity.

The present invention also provides methods of screening for modulators, e.g., activators, inhibitors, enhancers, etc., of mechanosensory transduction channels. Such modulators would be useful to alter mechanosensory transduction activity in an animal, e.g., for the treatment of any of a number of human disorders. Thus, the invention provides assays for mechanosensory transduction modulation, where the MSC proteins act as a direct or indirect reporter for mechanosensory transduction activity. MSC proteins can be used in assays, in vitro, in vivo, or ex vivo, to detect changes in ion flux, ion concentration, membrane potential, signal transduction, transcription, or other biological or biophysical effects of mechanical stimulus detection.

In one embodiment, MSC proteins can be used as indirect reporters via attachment to a second reporter molecule such as green fluorescent protein (see, e.g., Mistili & Spector, *Nature Biotechnology*, 15:961-964 (1997)). In one embodiment, MSC proteins are recombinantly expressed in cells, e.g., *Xenopus* oocytes, and modulation of mechanosensory transduction is assayed by detecting changes in transmembrane potential, mechanosensory potential, intracellular ion concentration, ion flux, and the like.

In certain embodiments, potential modulators are identified by virtue of an ability to physically interact with an MSC protein. Assays for physically-interacting molecules would provide an efficient primary screen for candidate MSC modulators, and, in addition, would allow the identification of proteins and other compounds that naturally interact with MSC proteins in vivo.

The invention also provides methods of detecting MSC nucleic acid and protein expression, allowing investigation into mechanosensory regulation and the identification of mechanosensory cells. The present nucleic acids and proteins can also be used to genotype an animal, including humans, for forensic, paternity, epidemiological, or other investigations. The present invention also provides conserved sequences found in multiple MSC sequences, allowing the identification of even distantly related MSC homologs (see, for example, SEQ ID NOs:7-9). In addition, the present invention provides methods for identifying mutations in a mechanosensory transduction channel protein that eliminate or reduce function of the channel. Such mutations likely underlie one or more of the human conditions involving loss of mechanosensation discussed herein. As such, the invention provides methods of diagnosing mechanosensory transduction defects in animals.

Functionally, the MSC proteins form, within a cell membrane, a channel that directly detects mechanical stimuli and, in response to the stimuli, allows the influx of cations into a cell, thereby depolarizing the cell and initiating an electrical, i.e. neural, signal.

Structurally, the nucleotide sequences of MSCs (see, e.g., SEQ ID NOs: 1, 3, and 5, representing the *Drosophila* genomic, *Drosophila* cDNA, and *Caenorhabditis elegans* genomic sequences, respectively) encode polypeptides of from about 1619-1709 amino acids with a predicted molecular weight of about 177 kDa (see, e.g., SEQ ID NOs:2, 4, and 6). The MSC genes typically contain about 19 exons, encoding a protein with about 27 ankyrin repeats and from 6-11, typically about 8, transmembrane domains. Such proteins are weakly related to the TRP family of epithelial cation channels. MSC homologs from other species typically share at least about 70% identity over a region of at least about 25 amino acids in length, preferably 50 to 100 amino acids in length.

The present invention provides nucleic acids comprising an MSC wherein the nucleic acid is less than 120, 90, 60, 30, 20, 10, or 7 kb. In addition, nucleic acids comprising MSCs are provided wherein the MSC polynucleotide is isolated away from its genomic neighbors, i.e., the nucleic acid does not comprise any genes that are located within the same genomic region as the MSC gene.

The present invention also provides polymorphic variants of the MSC depicted in SEQ ID NO:2: variant #1, in which an isoleucine residue is substituted for a leucine residue at amino acid position 6; variant #2, in which a glycine residue is substituted for an alanine residue at amino acid position 13; and variant #3, in which an arginine residue is substituted for a lysine residue at amino acid position 22.

The present invention also provides polymorphic variants of the MSC depicted in SEQ ID NO:4: variant #1, in which an isoleucine residue is substituted for a leucine residue at amino acid position 24; variant #2, in which an alanine residue is substituted for a glycine residue at amino acid position 26; and variant #3, in which an aspartic acid residue is substituted for a glutamic acid residue at amino acid position 30.

The present invention also provides mutated MSC sequences that eliminate mechanosensory transduction activity in vivo. For example, mutations that prematurely truncate MSC proteins in the ankyrin repeat region, or missense mutations that alter a cysteine residue between transmembrane segments four and five, e.g., a C to Y substitution, have been discovered that eliminate or severely reduce MSC activity. Such mutations can be used, e.g., to detect defects in mechanosensation, specifically in mechanosensory transduction channels, in an animal such as a human.

Specific regions of MSC may be used to identify polymorphic variants, interspecies homologs, and alleles of MSC. Such identification can be made in vitro, e.g., under stringent hybridization conditions or by PCR (e.g., using primers encoding SEQ ID NOs 7-9) and sequencing, or by using the sequence information provided herein in a computer system for comparison with other nucleotide sequences. Typically, identification of polymorphic variants and alleles of MSC proteins is made by comparing an amino acid sequence of about 25 amino acids or more, e.g., 50-100 amino acids. Amino acid identify of approximately at least about 70% or above, preferably 80%, most preferably 90-95% or above typically demonstrates that a protein is a polymorphic variant, interspecies homolog, or allele of MSC protein. Sequence comparison can be performed using any of the sequence comparison algorithms discussed herein. Antibodies that specifically bind to MSC protein or a conserved region thereof can also be used to identify alleles, interspecies homologs, and polymorphic variants.

Polymorphic variants, interspecies homologs, and alleles of MSC proteins can be confirmed by examining mechanosensory cell-specific expression of the putative MSC homolog. Typically, an MSC protein having a sequence of SEQ ID NO:2, 4, or 6 can be used as a positive control in comparison to the putative homolog. Such putative homologs are expected to retain the MSC structure described herein, i.e. intracellular domain with multiple, e.g., 27, ankyrin repeats, and a transmembrane domain containing multiple, e.g. 8, transmembrane domains.

The present invention also provides promoters, enhancers, 5'- and 3'-untranslated regions, and numerous other regulatory elements that control the transcription, translation, mRNA stability, mRNA localization, and other factors regulating MSC expression. For example, SEQ ID NO:1 provides genomic DNA sequence including MSC coding sequence as well as upstream and downstream regulatory sequences, including promoter sequences, etc. Promoters and other regulatory sequences can be identified using standard methods well known to those of skill in the art, including by homology to well conserved regulatory elements such as the TATA box or other elements, as taught, e.g., in Ausubel et al., supra, or in Lewin, *Genes IV* (1990). Promoter, enhancer, and other regulatory elements can also be determined functionally, e.g., by fusing specific regions of SEQ ID NO:1 to a reporter gene and determining which regions are sufficient for expression of the reporter gene, or by mutagenizing specific regions of SEQ ID NO:1 and thereby determining which regions are required for expression. Such methods are well known to those of skill in the art. Any of the present regulatory elements can be used in isolation or together, and can be used to drive the expression of an MSC protein, a marker protein, or any protein or RNA that is desirably expressed in a cell or other expression system. In preferred embodiments, an MSC regulatory element is used to drive the expression of a protein, e.g., an MSC or a heterologous polypeptide, in a tissue-specific manner, i.e., specifically in mechanosensory cells.

MSC nucleotide and amino acid sequences can also be used to construct models of mechanosensory transduction cell proteins in a computer system. Such models can be used, e.g., to identify compounds that may interact with, activate, or inhibit MSC protein channels. Such compounds can then be used for various applications, such as to modulate mechanosensory transduction activity in vivo or to investigate the various roles of MSC in mechanosensory transduction in vivo.

The isolation of MSC protein also provides a means for assaying for inhibitors and activators of mechanosensory

transduction channels, as well as for molecules, e.g., proteins, that interact with MSC proteins in vitro or in vivo. Biologically active MSC protein channels are useful for testing inhibitors and activators of MSC as mechanosensory transduction channels using in vivo and in vitro expression, e.g., in oocytes, and measuring MSC expression, phosphorylation state, membrane potential, mechanosensory potential, intra- or extra-cellular ion concentration, ion flux, and the like. Molecules can also be screened for the ability to physically interact with, e.g., bind to, MSC proteins, fragments thereof, or MSC nucleic acids, e.g., MSC promoter sequences, as shown in SEQ ID NO:1 and SEQ ID NO:3. Such interacting molecules can interact with any part of an MSC, e.g., the extracellular domain, transmembrane domain region, or intracellular domain, e.g., an ankyrin repeat. Such molecules may be involved in, or used to identify molecules capable of modulating, any aspect of MSC activity, including channel formation, detection of a mechanical stimulus, opening and/or closing of the channel, ion specificity of the channel, adaptation of the channel, or any other functional or physical aspect of the channel.

The present invention also provides assays, preferably high throughput assays, to identify molecules that interact with and/or modulate an MSC polypeptide. In numerous assays, a particular domain of an MSC is used, e.g., an extracellular, transmembrane, or intracellular domain. In numerous embodiments, an extracellular domain is bound to a solid substrate, and used, e.g., to isolate enhancers, inhibitors, or any molecule that can bind to and/or modulate the activity of an extracellular domain of an MSC polypeptide. In certain embodiments, a domain of an MSC polypeptide, e.g., an extracellular, transmembrane, or intracellular domain, is fused to a heterologous polypeptide, thereby forming a chimeric polypeptide. Such chimeric polypeptides are useful, e.g., in assays to identify modulators of an MSC polypeptide.

Such modulators and interacting molecules can be used for various purposes, such as to further investigate mechanosensory transduction channel activity in animal cells, or to modulate mechanosensory transduction activity in cells, e.g. to treat one or more conditions associated with a mechanosensory defect. It will be appreciated that in any of the binding assays or the in vitro or in vivo functional assays described herein, a full-length MSC can be used, or, alternatively, a fragment of an MSC can be used, for example a region containing only the ankyrin repeats, containing only the transmembrane domains, containing only the extracellular domain, or containing only a fragment of any these regions, will be used. Further, such fragments can be used alone, or fused to a heterologous protein any other molecule.

#### Definitions

The term "mechanosensory transduction protein" refers to a polypeptide that, when expressed in a cell or an oocyte, confers onto the cell an ability to detect changes in pressure, motion, or any other mechanical stimulus as described herein. Such proteins can be expressed naturally or recombinantly, and can confer such activity on the cell in vitro, in vivo, or ex vivo. Typically, such proteins will be at least about 70% identical to an amino acid sequence of SEQ ID NO:2, 4, or 6, and will include intracellular domains, including ankyrin repeats, and transmembrane domains. However, such proteins can also refer to one or more domains of these sequences in isolation, e.g., the ankyrin repeats, the extracellular domain, the transmembrane domains, or any subfragments thereof, alone. Such proteins can be involved in any mechanosensory process, such as tactile sensation, proprioception, hearing, baroreception, and others.



The term "MSC protein" refers to polymorphic variants, alleles, mutants, and interspecies homologs that: (1) have about 70% amino acid sequence identity, preferably about 85-90% amino acid sequence identity to SEQ ID NOS:2, 4, or 6 over a window of about 25 amino acids, preferably 50-100 amino acids; (2) bind to antibodies raised against an immunogen comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2, 4, 6-9, and conservatively modified variants thereof; (3) specifically hybridize (with a size of at least about 500, preferably at least about 900 nucleotides) under stringent hybridization and/or wash conditions to a sequence selected from the group consisting of SEQ ID NO:1, 3, and 5, and conservatively modified variants thereof; or (4) are amplified by primers that specifically hybridize under stringent hybridization conditions to the same sequence as a degenerate primer sets encoding SEQ ID NOS: 7-9.

"Biological sample" as used herein is a sample of biological tissue or fluid that contains an MSC protein or nucleic acid encoding an MSC protein. Such samples include, but are not limited to, tissue isolated from humans, mice, rats, and other animals. Biological samples may also include sections of tissues such as frozen sections taken for histological purposes. A biological sample is typically obtained from a eukaryotic organism, such as insects, protozoa, birds, fish, reptiles, and preferably a mammal such as rat, mouse, cow, dog, guinea pig, or rabbit, and most preferably a primate such as chimpanzees or humans. Preferred tissues include tissues involved in mechanosensation, such as the inner ear or any mechanosensory epithelial or neural tissue.

The phrase "functional effects" in the context of assays for testing compounds that modulate MSC protein-mediated mechanosensory transduction includes the determination of any parameter that is indirectly or directly under the influence of the channel. It includes changes in ion flux, membrane potential, current flow, transcription, MSC protein phosphorylation or dephosphorylation, signal transduction, in vitro, in vivo, and ex vivo and also includes other physiologic effects such increases or decreases of neurotransmitter or hormone release.

By "determining the functional effect" is meant assays for a compound that increases or decreases a parameter that is indirectly or directly under the influence of MSC proteins. Such functional effects can be measured by any means known to those skilled in the art, e.g., patch clamping, voltage-sensitive dyes, whole-cell currents, radioisotope efflux, inducible markers, oocyte MSC expression; tissue culture cell MSC expression; transcriptional activation of MSC protein; ligand-binding assays; membrane potential and conductance changes; ion-flux assays; changes in intracellular calcium levels; neurotransmitter release, and the like.

A "physical effect" in the context of assays for testing the ability of a compound to affect the activity of or bind to an MSC polypeptide refers to any detectable alteration in the physical property or behavior of an MSC polypeptide due to an interaction with a heterologous compound, or any detection of a physical interaction using, e.g., electrophoretic, chromatographic, or immunologically-based assay, or using a two-hybrid screen as described infra. For example, a physical effect can include any alteration in any biophysical property of an MSC channel comprising an MSC polypeptide, e.g., the cation specificity or mechanical sensitivity of the channel, or any structural or biochemical properties of an MSC polypeptide, e.g., its secondary, tertiary, or quaternary structure, hydrodynamic properties, spectral properties, chemical properties, or any other such property as described, e.g., in Creighton, *Proteins* (1984).

"Inhibitors," "activators," and "modulators" of MSC refer to any inhibitory or activating molecules identified using in vitro and in vivo assays for mechanosensory transduction, e.g., agonists, antagonists, and their homologs and mimetics. Inhibitors are compounds that decrease, block, prevent, delay activation, inactivate, desensitize, or down regulate mechanosensory transduction, e.g., antagonists. Activators are compounds that increase, open, activate, facilitate, enhance activation, sensitize or up-regulate mechanosensory transduction, e.g., agonists. Modulators include genetically-modified versions of MSC, e.g., with altered activity, as well as naturally-occurring and synthetic ligands, antagonists, agonists, small chemical molecules and the like. Such assays for inhibitors and activators include, e.g., expressing MSC protein in cells or cell membranes, applying putative modulator compounds, and then determining the functional effects on mechanosensory transduction, as described above. Samples or assays comprising MSC that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of inhibition. Control samples (untreated with inhibitors) are assigned a relative MSC activity value of 100%. Inhibition of MSC is achieved when the C activity value relative to the control is about 80%, preferably 50%, more preferably 25-1%. Activation of MSCs is achieved when the MSC activity value relative to the control is 110%, more preferably 150%, more preferably 200-500%, more preferably 1000-3000% higher.

"Biologically active" MSC refers to an MSC protein, or a nucleic acid encoding the MSC protein, having mechanosensory transduction activity as described above, involved in mechanosensory transduction in mechanosensory cells.

The terms "isolated" "purified" or "biologically pure" refer to material that is substantially or essentially free from components which normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high-performance liquid chromatography. A protein that is the predominant species present in a preparation is substantially purified. In particular, an isolated MSC nucleic acid is separated, e.g., from open reading frames or fragments of open reading frames, e.g., that naturally flank the MSC gene and encode proteins other than MSC protein. An isolated MSC nucleic acid is typically contiguous, i.e., heterologous sequences are typically not embedded in the MSC nucleic acid sequence, although heterologous sequences are often found adjoining an isolated MSC nucleic acid sequence. The term "purified" denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. Particularly, it means that the nucleic acid or protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure.

"Nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions)

and complementary sequences, as well as the sequence explicitly indicated. The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.

The terms "polypeptide," "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an analog or mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers. Polypeptides can be modified, e.g., by the addition of carbohydrate residues to form glycoproteins. The terms "polypeptide," "peptide" and "protein" include glycoproteins, as well as non-glycoproteins.

The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, i.e., a carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes (A, T, G, C, U, etc.).

"Conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the any position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues to yield a codon encoding the same amino acid residue (Batzer et al., *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608 (1985); Rossolini et al., *Mol. Cell. Probes* 8:91-98 (1994)). Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon in an amino acid herein, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to

yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants and alleles of the invention.

The following groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Glycine (G);
- 2) Serine (S), Threonine (T);
- 3) Aspartic acid (D), Glutamic acid (E);
- 4) Asparagine (N), Glutamine (Q);
- 5) Cysteine (C), Methionine (M);
- 6) Arginine (R), Lysine (K), Histidine (H);
- 7) Isoleucine (I), Leucine (L), Valine (V); and
- 8) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

(see, e.g., Creighton, *Proteins* (1984) for a discussion of amino acid properties).

A "label" or a "detectable moiety" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include <sup>32</sup>P, fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins which can be made detectable, e.g., by incorporating a radiolabel into the peptide or used to detect antibodies specifically reactive with the peptide.

A "labeled nucleic acid probe or oligonucleotide" is one that is bound, either covalently, through a linker or a chemical bond, or noncovalently, through ionic, van der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the probe may be detected by detecting the presence of the label bound to the probe.

As used herein, a "nucleic acid probe or oligonucleotide" is defined as a nucleic acid capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base pairing, usually through hydrogen bond formation. As used herein, a probe may include natural (i.e., A, G, C, or T) or modified bases (7-deazaguanosine, inosine, etc.). In addition, the bases in a probe may be joined by a linkage other than a phosphodiester bond, so long as it does not interfere with hybridization. Thus, for example, probes may be peptide nucleic acids in which the constituent bases are joined by peptide bonds rather than phosphodiester linkages. It will be understood by one of skill in the art that probes may bind target sequences lacking complete complementarity with the probe sequence depending upon the stringency of the hybridization conditions. The probes are preferably directly labeled as with isotopes, chromophores, lumiphores, chromogens, or indirectly labeled such as with biotin to which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the select sequence or subsequence.

The term "recombinant" when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is

derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

A "promoter" is defined as an array of nucleic acid control sequences that direct transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter, or array of transcription factor binding sites) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

An "expression vector" is a nucleic acid construct, generated recombinantly or synthetically, with a series of specified nucleic acid elements that permit transcription of a particular nucleic acid in a host cell. The expression vector can be part of a plasmid, virus, or nucleic acid fragment. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence over a comparison window, as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Such sequences are then said to be "substantially identical." This definition also refers to the complement of a test sequence. Preferably, the percent identity exists over a region of the sequence that is at least about 25 amino acids in length, more preferably over a region that is 50 or 100 amino acids in length.

For sequence comparison, one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to

600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by manual alignment and visual inspection.

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, e.g., version 7.0 (Devereaux et al., *Nuc. Acids Res.* 12:387-395 (1984)).

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment.

The BLAST program uses as defaults a wordlength (W) of 11, the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below.

The phrase "selectively (or specifically) hybridizes to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

The phrase "stringent hybridization conditions," or "stringent wash conditions," refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10° C. lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength pH. The  $T_m$  is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at  $T_m$ , 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, preferably 10 times background hybridization. Washes can be performed for varying amounts of time, e.g., 5 minutes, 15 minutes, 30 minutes, 1 hour or more. Exemplary stringent hybridization or wash conditions can be as following: 50% formamide,

5×SSC, and 1% SDS, incubating at 42° C., or, 5×SSC, 1% SDS, incubating at 65° C., with wash in 0.2×SSC, and 0.1% SDS at 65° C.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Exemplary "moderately stringent hybridization conditions," or "moderately stringent wash conditions," include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 1×SSC at 45° C. Washes can be performed for varying amounts of time, e.g., 5 minutes, 15 minutes, 30 minutes, 1 hour or more. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

A further indication that two polynucleotides are substantially identical is if the reference sequence, amplified by a pair of oligonucleotide primers, can then be used as a probe under stringent hybridization and/or wash conditions to isolate the test sequence from a cDNA or genomic library, or to identify the test sequence in, e.g., a northern or Southern blot. Alternatively, another indication that the sequences are substantially identical is if the same set of PCR primers can be used to amplify both sequences.

"Antibody" refers to a polypeptide substantially encoded by an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain ( $V_L$ ) and variable heavy chain ( $V_H$ ) refer to these light and heavy chains respectively.

Antibodies exist, e.g., as intact immunoglobulins or as a number of well characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce  $F(ab)_2$ , a dimer of Fab which itself is a light chain joined to  $V_H-C_H1$  by a disulfide bond. The  $F(ab)_2$  may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the  $F(ab)_2$  dimer into an Fab' monomer. The Fab' monomer is essentially an Fab with part of the hinge region (see *Fundamental Immunology* (Paul ed., 3d ed. 1993). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized de novo either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies or those synthesized de novo using recombinant DNA methodologies (e.g., single chain Fv).

A “chimeric antibody” is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, e.g., an enzyme, toxin, hormone, growth factor, drug, etc.; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

An “anti-*MSC*” antibody is an antibody or antibody fragment that specifically binds a polypeptide encoded by the *MSC* gene, cDNA, or a subsequence thereof.

The term “immunoassay” is an assay that uses an antibody to specifically bind an antigen. The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen.

The phrase “specifically (or selectively) binds” to an antibody or “specifically (or selectively) immunoreactive with,” when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to *MSC* protein from specific species such as rat, mouse, or human can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with *MSC* and not with other proteins, except for polymorphic variants and alleles of *MSC*. This selection may be achieved by subtracting out antibodies that cross-react with *MSC* proteins from other species. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (see, e.g., Harlow & Lane, *Antibodies, A Laboratory Manual* (1988), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

The phrase “selectively associates with” refers to the ability of a nucleic acid to “selectively hybridize” with another as defined above, or the ability of an antibody to “selectively (or specifically) bind to a protein, as defined above.

By “host cell” is meant a cell that contains an expression vector and supports the replication or expression of the expression vector. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells such as CHO, HeLa and the like, e.g., cultured cells, explants, and cells in vivo.

#### Isolation of *MSC* Nucleic Acids

##### General Recombinant DNA Methods

This invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include Sambrook et al., *Molecular Cloning, A Laboratory Manual* (2nd ed. 1989); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel et al., eds., 1994)).

For nucleic acids, sizes are given in either kilobases (kb) or base pairs (bp). These are estimates derived from agarose or

acrylamide gel electrophoresis, from sequenced nucleic acids, or from published DNA sequences. For proteins, sizes are given in kilodaltons (kDa) or amino acid residue numbers. Protein sizes are estimated from gel electrophoresis, from sequenced proteins, from derived amino acid sequences, or from published protein sequences.

Oligonucleotides that are not commercially available can be chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage & Caruthers, *Tetrahedron Letts.* 22:1859-1862 (1981), using an automated synthesizer, as described in Van Devanter et al., *Nucleic Acids Res.* 12:6159-6168 (1984). Purification of oligonucleotides is by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson & Reanier, *J. Chrom.* 255:137-149 (1983).

The sequence of the cloned genes and synthetic oligonucleotides can be verified after cloning using, e.g., the chain termination method for sequencing double-stranded templates of Wallace et al., *Gene* 16:21-26 (1981).

##### Cloning *MSC* Nucleic Acids

In general, the nucleic acid sequences encoding *MSC* and related nucleic acid sequence homologs are cloned from cDNA and genomic DNA libraries by hybridization with a probe, or isolated using amplification techniques with oligonucleotide primers. For example, *MSC* sequences are typically isolated from mammalian nucleic acid (genomic or cDNA) libraries by hybridizing with a nucleic acid probe, the sequence of which can be derived from SEQ ID NOS:1, 3, or 5. *MSC* RNA and cDNA can be isolated from any of a number of tissues, such as hair cells of the inner ear, sensory neurons, or any other mechanosensory cell.

Amplification techniques using primers can also be used to amplify and isolate an *MSC* polynucleotide from DNA or RNA. The degenerate primers encoding the following amino acid sequences can also be used to amplify a sequence of *MSC*: SEQ ID NOS:7-9 (see, e.g., Dieffenbach & Dveksler, *PCR Primer: A Laboratory Manual* (1995)). These primers can be used, e.g., to amplify either the full length sequence or a probe of one to several hundred nucleotides, which is then used to screen a mammalian library for full-length *MSC* sequences.

Nucleic acids encoding *MSC* proteins can also be isolated from expression libraries using antibodies as probes. Such polyclonal or monoclonal antibodies can be raised using polypeptides comprising the sequence of, e.g., SEQ ID NOS: 2, 4, 6, 7, 8 or 9.

##### cDNA and Genomic Libraries

*MSC* polymorphic variants, alleles, and interspecies homologs that are substantially identical to *MSC* proteins can be isolated using *MSC* nucleic acid probes, and oligonucleotides under stringent hybridization conditions, by screening libraries. Alternatively, expression libraries can be used to clone *MSC* and *MSC* polymorphic variants, alleles, and interspecies homologs, by detecting expressed homologs immunologically with antisera or purified antibodies made against *MSC*, which also recognize and selectively bind to the *MSC* homolog.

To make a cDNA library, one should choose a source that is rich in *MSC* mRNA, e.g., inner ear tissue or other sources of mechanosensory cells, e.g., sensory epithelial cells or neurons. The mRNA is then made into cDNA using reverse transcriptase, ligated into a recombinant vector, and transfected into a recombinant host for propagation, screening and cloning. Methods for making and screening cDNA libraries are well known (see, e.g., Gubler & Hoffman, *Gene* 25:263-269 (1983); Sambrook et al., *supra*; Ausubel et al., *supra*).

For a genomic library, the DNA is extracted from the tissue and either mechanically sheared or enzymatically digested to yield fragments of about 12-20 kb. The fragments are then separated by gradient centrifugation from undesired sizes and are constructed in bacteriophage lambda vectors. These vectors and phage are packaged in vitro. Recombinant phage are analyzed by plaque hybridization as described in Benton & Davis, *Science* 196:180-182 (1977). Colony hybridization is carried out as generally described in Grunstein et al., *Proc. Natl. Acad. Sci. USA.*, 72:3961-3965 (1975).

#### Amplification Methods

An alternative method of isolating MSC nucleic acid and its homologs combines the use of synthetic oligonucleotide primers and amplification of an RNA or DNA template (see U.S. Pat. Nos. 4,683,195 and 4,683,202; *PCR Protocols: A Guide to Methods and Applications* (Innis et al., eds, 1990)). Methods such as polymerase chain reaction (PCR) and ligase chain reaction (LCR) can be used to amplify nucleic acid sequences of MSC directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. Degenerate oligonucleotides can be designed to amplify MSC homologs using the sequences provided herein. Restriction endonuclease sites can be incorporated into the primers. Polymerase chain reaction or other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of MSC-encoding mRNA in physiological samples, for nucleic acid sequencing, or for other purposes. Genes amplified by the PCR reaction can be purified from agarose gels and cloned into an appropriate vector.

Gene expression of MSC protein can be analyzed by techniques known in the art, e.g., reverse transcription and amplification of mRNA, isolation of total RNA or poly A<sup>+</sup> RNA, Northern blotting, dot blotting, in situ hybridization, RNase protection, probing DNA microchip arrays, and the like. In one embodiment, high density oligonucleotide analysis technology (e.g., GeneChip™) is used to identify homologs and polymorphic variants of MSC. In the case where the homologs being identified are linked to a known disease, they can be used with GeneChip™ as a diagnostic tool in detecting the disease in a biological sample, see, e.g., Gunthand et al., *AIDS Res. Hum. Retroviruses* 14: 869-876 (1998); Kozal et al., *Nat. Med.* 2:753-759 (1996); Matson et al., *Anal. Biochem.* 224:110-106 (1995); Lockhart et al., *Nat. Biotechnol.* 14:1675-1680 (1996); Gingeras et al., *Genome Res.* 8:435-448 (1998); Hacia et al., *Nucleic Acids Res.* 26:3865-3866 (1998).

Synthetic oligonucleotides can be used to construct recombinant MSC genes for use as probes or for expression of protein. This method is performed using a series of overlapping oligonucleotides usually 40-120 bp in length, representing both the sense and nonsense strands of the gene. These DNA fragments are then annealed, ligated and cloned. Alternatively, amplification techniques can be used with precise primers to amplify a specific subsequence of the MSC nucleic acid. The specific subsequence is then ligated into an expression vector.

The nucleic acid encoding the MSC protein is typically cloned into intermediate vectors before transformation into prokaryotic or eukaryotic cells for replication and/or expression. These intermediate vectors are typically prokaryote vectors, e.g., plasmids, or shuttle vectors.

#### Expressing Nucleic Acids in Prokaryotes and Eukaryotes Expression Vectors

To obtain high level expression of a cloned gene or nucleic acid, such as those cDNAs encoding an MSC protein, one

typically subclones MSC into an expression vector that contains a strong promoter to direct transcription, a transcription/translation terminator, and if for a nucleic acid encoding a protein, a ribosome binding site for translational initiation. Suitable bacterial promoters are well known in the art and described, e.g., in Sambrook et al. and Ausubel et al. Bacterial expression systems for expressing the MSC protein are available in, e.g., *E. coli*, *Bacillus* sp., and *Salmonella* (Palva et al., *Gene* 22:229-235 (1983); Mosbach et al., *Nature* 302:543-545 (1983)). Kits for such expression systems are commercially available. Eukaryotic expression systems for mammalian cells, yeast, and insect cells are well known in the art and are also commercially available.

#### Promoters

The promoter used to direct expression of a heterologous nucleic acid depends on the particular application. The promoter is preferably positioned about the same distance from the heterologous transcription start site as it is from the transcription start site in its natural setting. As is known in the art, however, some variation in this distance can be accommodated without loss of promoter function.

In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the MSC-encoding nucleic acid in host cells. A typical expression cassette thus contains a promoter operably linked to the nucleic acid sequence encoding MSC protein and signals required for efficient polyadenylation of the transcript, ribosome binding sites, and translation termination. The nucleic acid sequence encoding MSC protein may typically be linked to a cleavable signal peptide sequence to promote secretion of the encoded protein by the transformed cell. Such signal peptides would include, among others, the signal peptides from tissue plasminogen activator, insulin, and neuron growth factor, and juvenile hormone esterase of *Heliothis virescens*. Additional elements of the cassette may include enhancers and, if genomic DNA is used as the structural gene, introns with functional splice donor and acceptor sites.

#### Other Elements

In addition to a promoter sequence, the expression cassette should also contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

The particular expression vector used to transport the genetic information into the cell is not particularly critical. Any of the conventional vectors used for expression in eukaryotic or prokaryotic cells may be used. Standard bacterial expression vectors include plasmids such as pBR322 based plasmids, pSKF, pET23D, and fusion expression systems such as GST and LacZ. Epitope tags can also be added to recombinant proteins to provide convenient methods of isolation, e.g., c-myc.

Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A<sup>+</sup>, pMTO10/A<sup>+</sup>, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

Some expression systems have markers that provide gene amplification such as thymidine kinase, hygromycin B phos-

phototransferase, and dihydrofolate reductase. Alternatively, high yield expression systems not involving gene amplification are also suitable, such as using a baculovirus vector in insect cells, with a MSC encoding sequence under the direction of the polyhedrin promoter or other strong baculovirus promoters.

The elements that are typically included in expression vectors also include a replicon that functions in *E. coli*, a gene encoding antibiotic resistance to permit selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of eukaryotic sequences. The particular antibiotic resistance gene chosen is not critical, any of the many resistance genes known in the art are suitable. The prokaryotic sequences are preferably chosen such that they do not interfere with the replication of the DNA in eukaryotic cells, if necessary.

#### Transfection Methods

Standard transfection methods are used to produce bacterial, mammalian, yeast or insect cell lines that express large quantities of MSC protein, which are then purified using standard techniques (see, e.g., Colley et al., *J. Biol. Chem.* 264:17619-17622 (1989); *Guide to Protein Purification, in Methods in Enzymology*, vol. 182 (Deutscher, ed., 1990)). Transformation of eukaryotic and prokaryotic cells are performed according to standard techniques (see, e.g., Morrison, *J. Bact.* 132:349-351 (1977); Clark-Curtiss & Curtiss, *Methods in Enzymology* 101:347-362 (Wu et al., eds, 1983)).

Any of the well known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (see, e.g., Sambrook et al., supra). It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing MSC.

After the expression vector is introduced into the cells, the transfected cells are cultured under conditions favoring expression of MSC, which is recovered from the culture using standard techniques identified below.

#### Purification of MSC Proteins

Either naturally occurring or recombinant MSC protein can be purified for use in functional assays. Preferably, recombinant MSC is purified. Naturally occurring MSC is purified, e.g., from mammalian tissue such as inner ear tissue or other tissues including mechanosensory cells. Recombinant MSC is purified from any suitable expression system.

MSC protein may be purified to substantial purity by standard techniques, including selective precipitation with such substances as ammonium sulfate; column chromatography, immunopurification methods, and others (see, e.g., Scopes, *Protein Purification: Principles and Practice* (1982); U.S. Pat. No. 4,673,641; Ausubel et al., supra; and Sambrook et al., supra).

A number of procedures can be employed when recombinant MSC is being purified. For example, proteins having established molecular adhesion properties can be reversibly fused to MSC. With the appropriate ligand, MSC can be selectively adsorbed to a purification column and then freed from the column in a relatively pure form. The fused protein is then removed by enzymatic activity. Finally MSC could be purified using immunoaffinity columns.

#### Purification from Recombinant Bacteria

Recombinant proteins are expressed by transformed bacteria in large amounts, typically after promoter induction; but expression can be constitutive. Promoter induction with IPTG is one example of an inducible promoter system. Bacteria are grown according to standard procedures in the art. Fresh or frozen bacteria cells are used for isolation of protein.

Proteins expressed in bacteria may form insoluble aggregates ("inclusion bodies"). Several protocols are suitable for purification of MSC inclusion bodies. For example, purification of inclusion bodies typically involves the extraction, separation and/or purification of inclusion bodies by disruption of bacterial cells, e.g., by incubation in a buffer of 50 mM TRIS/HCL pH 7.5, 50 mM NaCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 0.1 mM ATP, and 1 mM PMSF. The cell suspension can be lysed using 2-3 passages through a French Press, homogenized using a Polytron (Brinkman Instruments) or sonicated on ice. Alternate methods of lysing bacteria are apparent to those of skill in the art (see, e.g., Sambrook et al., supra; Ausubel et al., supra).

If necessary, the inclusion bodies are solubilized, and the lysed cell suspension is typically centrifuged to remove unwanted insoluble matter. Proteins that formed the inclusion bodies may be renatured by dilution or dialysis with a compatible buffer. Suitable solvents include, but are not limited to urea (from about 4 M to about 8 M), formamide (at least about 80%, volume/volume basis), and guanidine hydrochloride (from about 4 M to about 8 M). Some solvents which are capable of solubilizing aggregate-forming proteins, for example SDS (sodium dodecyl sulfate), 70% formic acid, are inappropriate for use in this procedure due to the possibility of irreversible denaturation of the proteins, accompanied by a lack of immunogenicity and/or activity. Although guanidine hydrochloride and similar agents are denaturants, this denaturation is not irreversible and renaturation may occur upon removal (by dialysis, for example) or dilution of the denaturant, allowing re-formation of immunologically and/or biologically active protein. Other suitable buffers are known to those skilled in the art. MSC is separated from other bacterial proteins by standard separation techniques, e.g., with Ni-NTA agarose resin.

Alternatively, it is possible to purify MSC protein from bacteria periplasm. After lysis of the bacteria, when MSC is exported into the periplasm of the bacteria, the periplasmic fraction of the bacteria can be isolated by cold osmotic shock in addition to other methods known to skill in the art. To isolate recombinant proteins from the periplasm, the bacterial cells are centrifuged to form a pellet. The pellet is resuspended in a buffer containing 20% sucrose. To lyse the cells, the bacteria are centrifuged and the pellet is resuspended in ice-cold 5 mM MgSO<sub>4</sub> and kept in an ice bath for approximately 10 minutes. The cell suspension is centrifuged and the supernatant decanted and saved. The recombinant proteins present in the supernatant can be separated from the host proteins by standard separation techniques well known to those of skill in the art.

#### Standard Protein Purification Techniques

##### Solubility Fractionation

Often as an initial step, particularly if the protein mixture is complex, an initial salt fractionation can separate many of the unwanted host cell proteins (or proteins derived from the cell culture media) from the recombinant protein of interest. The preferred salt is ammonium sulfate. Ammonium sulfate precipitates proteins by effectively reducing the amount of water in the protein mixture. Proteins then precipitate on the basis of their solubility. The more hydrophobic a protein is, the more likely it is to precipitate at lower ammonium sulfate concen-



trations. A typical protocol includes adding saturated ammonium sulfate to a protein solution so that the resultant ammonium sulfate concentration is between 20-30%. This concentration will precipitate the most hydrophobic of proteins. The precipitate is then discarded (unless the protein of interest is hydrophobic) and ammonium sulfate is added to the supernatant to a concentration known to precipitate the protein of interest. The precipitate is then solubilized in buffer and the excess salt removed if necessary, either through dialysis or diafiltration. Other methods that rely on solubility of proteins, such as cold ethanol precipitation, are well known to those of skill in the art and can be used to fractionate complex protein mixtures.

#### Size Differential Filtration

The molecular weight of MSC protein can be used to isolate it from proteins of greater and lesser size using ultrafiltration through membranes of different pore size (for example, Amicon or Millipore membranes). As a first step, the protein mixture is ultrafiltered through a membrane with a pore size that has a lower molecular weight cut-off than the molecular weight of the protein of interest. The retentate of the ultrafiltration is then ultrafiltered against a membrane with a molecular cut off greater than the molecular weight of the protein of interest. The recombinant protein will pass through the membrane into the filtrate. The filtrate can then be chromatographed as described below.

#### Column Chromatography

MSC proteins can also be separated from other proteins on the basis of its size, net surface charge, hydrophobicity, and affinity for ligands. In addition, antibodies raised against proteins can be conjugated to column matrices and the proteins immunopurified. All of these methods are well known in the art. It will be apparent to one of skill that chromatographic techniques can be performed at any scale and using equipment from many different manufacturers (e.g., Pharmacia Biotech).

#### Affinity-Based Techniques

Any of a number of affinity based techniques can be used to isolate MSC proteins from cells, cell extracts, or other sources. For example, affinity columns can be made using anti-MSC antibodies or other MSC-binding proteins, or physically-interacting proteins can be identified by co-immunoprecipitation or other methods. Such methods are well known to those of skill in the art and are taught, e.g., in Ausubel et al., Sambrook et al., Harlow and Lane, all supra.

#### Immunological Detection

In addition to the detection of MS genes and gene expression using nucleic acid hybridization technology, one can also use immunoassays to detect MSC proteins, e.g., to identify mechanosensory cells and variants of MSC proteins. Immunoassays can be used to qualitatively or quantitatively analyze MSC proteins. A general overview of the applicable technology can be found in Harlow & Lane, *Antibodies: A Laboratory Manual* (1988).

#### Antibodies to MSC Proteins

Methods of producing polyclonal and monoclonal antibodies that react specifically with MSC proteins are known to those of skill in the art (see, e.g., Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, supra; Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986); and Kohler & Milstein, *Nature* 256:495-497 (1975)). Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors, as well as preparation of polyclonal and

monoclonal antibodies by immunizing rabbits or mice (see, e.g., Huse et al., *Science* 246:1275-1281 (1989); Ward et al., *Nature* 341:544-546 (1989)).

A number of MSC peptides or a full-length protein may be used to produce antibodies specifically reactive with MSC protein. For example, recombinant MSC protein, or an antigenic fragment thereof, is isolated as described herein. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described above, and purified as generally described above. Recombinant protein is the preferred immunogen for the production of monoclonal or polyclonal antibodies. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Naturally occurring protein may also be used either in pure or impure form. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated, for subsequent use in immunoassays to measure the protein.

Methods of production of polyclonal antibodies are known to those of skill in the art. An inbred strain of mice (e.g., BALB/C mice) or rabbits is immunized with the protein using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to MSC proteins. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired (see, Harlow & Lane, supra).

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (see, Kohler & Milstein, *Eur. J. Immunol.* 6:511-519 (1976)). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse et al., *Science* 246:1275-1281 (1989).

Monoclonal antibodies and polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Typically, polyclonal antisera with a titer of  $10^4$  or greater are selected and tested for their cross reactivity against non-MSC proteins or even other related proteins from other organisms, using a competitive binding immunoassay. Specific polyclonal antisera and monoclonal antibodies will usually bind with a  $K_d$  of at least about 0.1 mM, more usually at least about 1  $\mu$ M, preferably at least about 0.1  $\mu$ M or better, and most preferably, 0.01  $\mu$ M or better.

Once MSC specific antibodies are available, MSC proteins can be detected by a variety of immunoassay methods. For a review of immunological and immunoassay procedures, see *Basic and Clinical Immunology* (Stites & Terr eds., 7th ed. 1991). Moreover, the immunoassays of the present invention can be performed in any of several configurations, which are



reviewed extensively in *Enzyme Immunoassay* (Maggio, ed., 1980); and Harlow & Lane, supra.

#### Immunological Binding Assays

MSC proteins can be detected and/or quantified using any of a number of well recognized immunological binding assays (see, e.g., U.S. Pat. Nos. 4,366,241, 4,376,110; 4,517, 288; and 4,837,168). For a review of the general immunoassays, see also *Methods in Cell Biology: Antibodies in Cell Biology*, volume 37 (Asai, ed. 1993); *Basic and Clinical Immunology* (Stites & Terr, eds., 7th ed. 1991). Immunological binding assays (or immunoassays) typically use an antibody that specifically binds to a protein or antigen of choice (in this case the MSC protein or antigenic subsequence thereof). The antibody (e.g., anti-MSC) may be produced by any of a number of means well known to those of skill in the art and as described above.

Immunoassays also often use a labeling agent to specifically bind to and label the complex formed by the antibody and antigen. The labeling agent may itself be one of the moieties comprising the antibody/antigen complex. Thus, the labeling agent may be a labeled MSC polypeptide or a labeled anti-MSC antibody. Alternatively, the labeling agent may be a third moiety, such as a secondary antibody, that specifically binds to the antibody/MSC complex (a secondary antibody is typically specific to antibodies of the species from which the first antibody is derived). Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (see, e.g., Kronval et al., *J. Immunol.* 111:1401-1406 (1973); Akerstrom et al., *J. Immunol.* 135:2589-2542 (1985)). The labeling agent can be modified with a detectable moiety, such as biotin, to which another molecule can specifically bind, such as streptavidin. A variety of detectable moieties are well known to those skilled in the art.

Throughout the assays, incubation and/or washing steps may be required after each combination of reagents. Incubation steps can vary from about 5 seconds to several hours, preferably from about 5 minutes to about 24 hours. However, the incubation time will depend upon the assay format, antigen, volume of solution, concentrations, and the like. Usually, the assays will be carried out at ambient temperature, although they can be conducted over a range of temperatures, such as 11° C. to 40° C.

#### Non-Competitive Formats

Immunoassays for detecting MSC proteins in samples may be either competitive or noncompetitive. Noncompetitive immunoassays are assays in which the amount of antigen is directly measured. In one preferred "sandwich" assay, for example, the anti-MSC antibodies can be bound directly to a solid substrate on which they are immobilized. These immobilized antibodies then capture MSC proteins present in the test sample. The MSC protein is thus immobilized and then bound by a labeling agent, such as a second MSC antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second or third antibody is typically modified with a detectable moiety, such as biotin, to which another molecule specifically binds, e.g., streptavidin, to provide a detectable moiety.

#### Competitive Formats

In competitive assays, the amount of MSC proteins present in the sample is measured indirectly by measuring the amount of a known, added (exogenous) MSC proteins displaced (competed away) from an anti-MSC antibody by the

unknown MSC protein present in a sample. In one competitive assay, a known amount of MSC protein is added to a sample and the sample is then contacted with an antibody that specifically binds to MSC proteins. The amount of exogenous MSC protein bound to the antibody is inversely proportional to the concentration of MSC protein present in the sample. In a particularly preferred embodiment, the antibody is immobilized on a solid substrate. The amount of MSC protein bound to the antibody may be determined either by measuring the amount of MSC protein present in a MSC protein/antibody complex, or alternatively by measuring the amount of remaining uncomplexed protein. The amount of MSC protein may be detected by providing a labeled MSC protein molecule.

A hapten inhibition assay is another preferred competitive assay. In this assay the known MSC protein, is immobilized on a solid substrate. A known amount of anti-MSC antibody is added to the sample, and the sample is then contacted with the immobilized MSC protein. The amount of anti-MSC antibody bound to the known immobilized MSC protein is inversely proportional to the amount of MSC protein present in the sample. Again, the amount of immobilized antibody may be detected by detecting either the immobilized fraction of antibody or the fraction of the antibody that remains in solution. Detection may be direct where the antibody is labeled or indirect by the subsequent addition of a labeled moiety that specifically binds to the antibody as described above.

#### Cross-Reactivity Determination

Immunoassays in the competitive binding format can also be used for crossreactivity determinations. For example, a protein at least partially encoded by SEQ ID NOS:1, 3, or 5 can be immobilized to a solid support. Proteins (e.g., MSC proteins and homologs) are added to the assay that compete for binding of the antisera to the immobilized antigen. The ability of the added proteins to compete for binding of the antisera to the immobilized protein is compared to the ability of MSC protein encoded by SEQ ID NO:1, 3, or 5 to compete with itself. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the added proteins listed above are selected and pooled. The cross-reacting antibodies are optionally removed from the pooled antisera by immunoabsorption with the added considered proteins, e.g., distantly related homologs. In one embodiment, antibodies that crossreact with MSC proteins from a different species are selectively removed, thereby enhancing the species-specificity of the antisera. For example, to obtain antibodies that specifically react with *Drosophila* MSC, the ability of SEQ ID NO:4 and SEQ ID NO:6 to compete for binding to antisera directed against SEQ ID NO:4 are compared, and antibodies that cross-react with SEQ ID NO:6 selectively removed.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein, thought to be perhaps an allele or polymorphic variant of MSC protein, to the immunogen protein (i.e., MSC protein of SEQ ID NOS:2, 4, 6-9). In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required to inhibit 50% of binding is less than 10 times the amount of the protein encoded by SEQ ID NOS:1, 3, or 5 that is required to inhibit 50% of binding, then the second protein is said to specifically bind to the polyclonal antibodies generated to an MSC protein immunogen.

## Other Formats

Western blot (immunoblot) analysis is used to detect and quantify the presence of MSC protein in the sample. The technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with the antibodies that specifically bind MSC protein. The anti-MSC antibodies specifically bind to the MSC protein on the solid support. These antibodies may be directly labeled or alternatively may be subsequently detected using labeled antibodies (e.g., labeled sheep anti-mouse antibodies) that specifically bind to the anti-MSC antibodies.

Other assay formats include liposome immunoassays (LIA), which use liposomes designed to bind specific molecules (e.g., antibodies) and release encapsulated reagents or markers. The released chemicals are then detected according to standard techniques (see, Monroe et al., *Amer. Clin. Prod. Rev.* 5:34-41 (1986)).

## Reduction of Non-Specific Binding

One of skill in the art will appreciate that it is often desirable to minimize non-specific binding in immunoassays. Particularly where the assay involves an antigen or antibody immobilized on a solid substrate, it is desirable to minimize the amount of non-specific binding to the substrate. Means of reducing such non-specific binding are well known to those of skill in the art. Typically, this technique involves coating the substrate with a proteinaceous composition. In particular, protein compositions such as bovine serum albumin (BSA), nonfat powdered milk, and gelatin are widely used, with powdered milk being most preferred.

## Labels

The particular label or detectable group used in the assay is not a critical aspect of the invention, as long as it does not significantly interfere with the specific binding of the antibody used in the assay. The detectable group can be any material having a detectable physical or chemical property. Such detectable labels have been well-developed in the field of immunoassays and, in general, most any label useful in such methods can be applied to the present invention. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include magnetic beads (e.g., DYNABEADS™), fluorescent dyes (e.g., fluorescein isothiocyanate, Texas red, rhodamine, and the like), radiolabels (e.g., <sup>3</sup>H, <sup>125</sup>I, <sup>35</sup>S, <sup>14</sup>C, or <sup>32</sup>P), enzymes (e.g., horse radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold or colored glass or plastic beads (e.g., polystyrene, polypropylene, latex, etc.).

The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide variety of labels may be used, with the choice of label depending on sensitivity required, ease of conjugation with the compound, stability requirements, available instrumentation, and disposal provisions.

Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (e.g., biotin) is covalently bound to the molecule. The ligand then binds to another molecule (e.g., streptavidin) molecule, which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. The ligands and their targets can be used in any suitable combination with

antibodies that recognize MSC protein, or secondary antibodies that recognize anti-MSC protein.

The molecules can also be conjugated directly to signal generating compounds, e.g., by conjugation with an enzyme or fluorophore. Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and glycosidases, or oxidotases, particularly peroxidases. Fluorescent compounds include fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, etc. Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, e.g., luminol. For a review of various labeling or signal producing systems that may be used, see U.S. Pat. No. 4,391,904.

Means of detecting labels are well known to those of skill in the art. Thus, for example, where the label is a radioactive label, means for detection include a scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence. The fluorescence may be detected visually, by means of photographic film, by the use of electronic detectors such as charge coupled devices (CCDs) or photomultipliers and the like. Similarly, enzymatic labels may be detected by providing the appropriate substrates for the enzyme and detecting the resulting reaction product. Finally simple calorimetric labels may be detected simply by observing the color associated with the label. Thus, in various dipstick assays, conjugated gold often appears pink, while various conjugated beads appear the color of the bead.

Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies. In this case, antigen-coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need be labeled and the presence of the target antibody is detected by simple visual inspection.

## Assays for Modulators of Mechanosensory Transduction

In numerous embodiments of this invention, assays will be performed to detect compounds that affect mechanosensory transduction in a cell. Such assays can involve the identification of compounds that interact with MSC proteins, either physically or genetically, and can thus rely on any of a number of standard methods to detect physical or genetic interactions between compounds. Such assays can also involve the detection of mechanosensory transduction in a cell or cell membrane, either in vitro or in vivo, and can thus involve the detection of transduction activity in the cell through any standard assay, e.g., by measuring ion flux, changes in membrane potential, and the like. Such cell-based assays can be performed in any type of cell, e.g., a sensory cell that naturally expresses MSC, a cultured cell that produces MSC due to recombinant expression, or, preferably, an oocyte that is induced to produce MSC through any of a number of means, as described infra.

In any of the binding or functional assays described herein, in vivo or in vitro, any MSC protein, or any derivative, variation, homolog, or fragment of an MSC protein, can be used. Preferably, the MSC protein is at least about 70% identical to SEQ ID NO:2, 4, or 6, and/or comprises SEQ ID NO:7, 8, or 9. In numerous embodiments, a fragment of an MSC protein is used. For example, a fragment that contains only the extracellular region, the ankyrin repeat region, or the transmembrane domains, i.e. the channel region (see, e.g., SEQ ID NOs:10-17), can be used. Such fragments can be used alone, in combination with other MSC fragments, or in combination

with sequences from a heterologous protein, e.g., the fragments can be fused to a heterologous polypeptide, thereby forming a chimeric polypeptide. Any individual domain or sequence, however small, can readily be used in the present invention, e.g., a single ankyrin repeat, transmembrane domain, etc., alone or in combination with other domains or with sequences from heterologous proteins. Such fragments and isolated domains of MSC proteins comprise an essential aspect of the present invention, and are of substantial importance in the assays described herein.

#### Assays for MSC-Interacting Compounds

In certain embodiments, assays will be performed to identify molecules that physically or genetically interact with MSC proteins. Such molecules can be any type of molecule, including polypeptides, polynucleotides, amino acids, nucleotides, carbohydrates, lipids, or any other organic or inorganic molecule. Such molecules may represent molecules that normally interact with MSC to effect mechanosensation in sensory cells, or may be synthetic or other molecules that are capable of interacting with MSC and which can potentially be used to modulate MSC activity in cells, or used as lead compounds to identify classes of molecules that can interact with and/or modulate MSC. Such assays may represent physical binding assays, such as affinity chromatography, immunoprecipitation, two-hybrid screens, or other binding assays, or may represent genetic assays as described infra.

Such interacting molecules may interact with any part of an MSC protein, e.g., the extracellular domain, the transmembrane domain region, or the intracellular domain, including the ankyrin repeats. MSC proteins act in sensory cells to depolarize the cell in response to a mechanical input outside of the cell. As such, interacting molecules may include those that interact with the extracellular domain of the protein, and which may enhance, inhibit, or otherwise modulate the detection of a mechanical input, and which may be part of, or interact with, an extracellular structure involved in mechanical detection, such as the stereocilium of a hair cell. An interacting molecule may also interact with the transmembrane domain region of the protein, and may be involved in, or capable of modulating, the formation of a channel, the opening or closing of a channel, etc. In addition, an interacting molecule may interact with an intracellular part of a channel, e.g., an ankyrin repeat, and be involved in, e.g., the function, regulation, adaptation, or any other aspect of channel activity.

The MSC protein used in such assays can be a full-length MSC protein or any subdomain of an MSC protein. In preferred embodiments, a fragment of an MSC protein comprising an extracellular domain of an MSC will be used. Molecules that bind to the extracellular domain of an MSC are particularly useful for the identification of modulators of MSC activity, as they are typically soluble and readily included in high throughput screening assay formats, as described infra.

#### Assays for Physical Interactions

Compounds that interact with MSC proteins can be isolated based on an ability to specifically bind to an MSC protein or fragment thereof. In numerous embodiments, the MSC protein or protein fragment will be attached to a solid support. In one embodiment, affinity columns are made using the MSC polypeptide, and physically-interacting molecules are identified. It will be apparent to one of skill that chromatographic techniques can be performed at any scale and using equipment from many different manufacturers (e.g., Pharmacia Biotech). In addition, molecules that interact with MSC proteins *in vivo* can be identified by co-immunoprecipitation or other methods, i.e. immunoprecipitating MSC proteins using anti-MSC antibodies from a cell or cell extract, and

identifying compounds, e.g., proteins, that are precipitated along with the MSC protein. Such methods are well known to those of skill in the art and are taught, e.g., in Ausubel et al., Sambrook et al., Harlow & Lane, all supra.

Two-hybrid screens can also be used to identify polypeptides that interact *in vivo* with an MSC or a fragment thereof (Fields et al., *Nature* 340:245-246 (1989)). Such screens comprise two discrete, modular domains of a transcriptional activation domain, which are produced in a cell as two separate polypeptides, each of which also comprises one of two potentially binding polypeptides. If the two potentially binding polypeptides in fact interact *in vivo*, then the DNA binding and the transcriptional activating domain of the transcription factor are united, thereby producing expression of a target gene in the cell. The target gene typically encodes an easily detectable gene product, e.g.,  $\beta$ -galactosidase, which can be detected using standard methods. In the present invention, an MSC polypeptide is fused to one of the two domains of the transcription factor, and the potential MSC-binding polypeptides (e.g., encoded by a cDNA library) are fused to the other domain. Such methods are well known to those of skill in the art, and are taught, e.g., in Ausubel et al., supra.

#### Assays for Genetic Interactions

It is expected that MSCs are assembled into multi-protein complexes in which the interactions are mediated by the large number of ankyrin repeats found in the N terminus of the protein. Genetic screens can thus be performed to identify such additional proteins that are involved in the transduction pathway. For example, genetic strains are produced that possess only a partially functional *nompC* (MSC) gene, which confers an incomplete mechanical sensitivity to the fly. Ideally, a vial of these flies would produce only 10-20 viable homozygotes. In this sensitized genetic background, flies will be screened for mutations in other genes that either suppress or enhance the survival of the mutant flies. Flies will be mutagenized using any standard chemical, radiation-based, or genetic method and then crossed into the above-described sensitized genetic background, followed by counting the number of homozygous progeny. Mutations that produce more than 10-20 flies per vial are considered suppressors of *nompC*, and those that produce fewer flies are considered enhancers. Similar screens can be performed using MSC genes in genetically tractable mammals, e.g., mice.

#### Assays for MSC Activity

The activity of MSC polypeptides, and any homolog, variant, derivative, or fragment thereof can be assessed using a variety of *in vitro* and *in vivo* assays for mechanoreceptor potential, e.g., measuring current, measuring membrane potential, measuring ion flux, e.g., potassium or calcium, measuring transcription levels, measuring neurotransmitter levels, using e.g., voltage-sensitive dyes, radioactive tracers, patch-clamp electrophysiology, transcription assays, and the like. Furthermore, such assays can be used to test for modulators, e.g., inhibitors or activators, of MSC. Such modulators can be a protein, amino acid, nucleic acid, nucleotide, lipid, carbohydrate, or any type of organic or inorganic molecule, including genetically altered versions of MSC proteins. Such assays can be performed using any of a large number of cells, including oocytes, cultured cells, sensory epithelial or neural cells, and others, and can be present *in vitro* or *in vivo*. Such cells can contain naturally expressed MSC, can be induced to express MSC using recombinant or other methods, or can comprise MSC by direct addition of the protein to the cell or cell membrane. In numerous embodiments, the cell or cell membrane comprising the MSC polypeptide will be anchored to a solid support.

Preferably, the MSC proteins used in the assay is selected from a polypeptide having a sequence of SEQ ID NOS:2, 4, or 6, or a conservatively modified variant thereof. Alternatively, the MSC protein used in the assay will be derived from a eukaryote and include an amino acid subsequence having amino acid sequence identity SEQ ID NOS:2, 4, or 6. Generally, the amino acid sequence identity will be at least 70%, preferably at least 85%, most preferably at least 90-95%. In preferred embodiments, a polypeptide comprising an extracellular domain is used, e.g., an extracellular domain of SEQ ID NO:2, 4, or 6. In such embodiments, the extracellular domain is often fused to a heterologous polypeptide, forming a chimeric polypeptide. Typically, such chimeric polypeptides will comprise an extracellular domain as well as multiple transmembrane domains, and will have mechanosensory transduction activity.

#### Detecting Mechanosensory Transduction

In numerous embodiments of the present invention, assays will be performed to detect alterations in an MSC protein, e.g., one expressed in a cell or cell membrane, or in mechanosensory transduction, or mechanoreceptor potential, in a cell or cell membrane, e.g., as a result of a mutation in an MSC or due to the presence of an MSC-modulating compound. Mechanosensory transduction or mechanoreceptor potential can be detected in any of a number of ways, including by detecting changes in ion flux, changes in polarization of a cell or cell membrane, changes in current, and other methods, including by measuring downstream cellular effects, e.g., neuronal signaling.

Changes in ion flux may be assessed by determining changes in polarization (i.e., electrical potential) of the cell or membrane expressing MSC. One means to determine changes in cellular polarization is by measuring changes in current (thereby measuring changes in polarization) with voltage-clamp and patch-clamp techniques, e.g., the "cell-attached" mode, the "inside-out" mode, and the "whole cell" mode (see, e.g., Ackerman et al., *New Engl. J. Med.* 336: 1575-1595 (1997)). Whole cell currents are conveniently determined using the standard methodology (see, e.g., Hamil et al., *Pflugers. Archiv.* 391:85 (1981)). Other known assays include: radioactive ion flux assays and fluorescence assays using voltage-sensitive dyes (see, e.g., Vestergarrd-Bogind et al., *J. Membrane Biol.* 88:67-75 (1988); Gonzales & Tsien, *Chem. Biol.* 4:269-277 (1997); Daniel et al., *J. Pharmacol. Meth.* 25:185-193 (1991); Holevinsky et al., *J. Membrane Biology* 137:59-70 (1994)). Generally, candidate compounds are tested in the range from 1 pM to 100 mM.

The effects of the test compounds, or sequence variation, upon the function of the MSC polypeptides can be measured by examining any of the parameters described above. In addition, any suitable physiological change that affects MSC activity, or reflects MSC activity, can be used to assess the influence of a test compound or sequence alteration on the MSC polypeptides of this invention. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as transmitter release, hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., northern blots), changes in cell metabolism such as cell growth or pH changes, and other effects.

Preferred assays for mechanosensory transduction channels include cells, e.g., oocytes, that are loaded with ion or voltage sensitive dyes to report receptor activity. Assays for determining activity of such receptors can also use known agonists and antagonists for other cation channels as negative or positive controls to assess activity of tested compounds. In assays for identifying modulatory compounds (e.g., agonists,

antagonists), changes in the level of ions in the cytoplasm or membrane voltage will be monitored using an ion-sensitive or membrane voltage fluorescent indicator, respectively. Among the ion-sensitive indicators and voltage probes that may be employed are those disclosed in the Molecular Probes 1997 Catalog. In addition, changes in cytoplasmic calcium, potassium, or other ion levels can be used to assess MSC function.

#### In vivo Assays

In certain embodiments, the mechanosensory activity of a cell will be examined in vivo. Such embodiments are useful for, e.g., examining the activity of an MSC or an MSC mutant, derivative, homolog, fragment, etc. Also, such assays are useful for detecting the activity of candidate MSC modulator in vivo. Potential MSCs can be produced in transgenic flies carrying the candidate cDNA driven by a suitable, e.g. a *nompC*, promoter/enhancer construct. These candidate channels can be expressed in mechanosensory neurons of flies and their mechano-electrical activity measured with bristle recordings. Methods of producing transgenic flies and methods of detecting mechanosensory transduction activity in fly mechanosensory neurons are well known to those of skill in the art and are described, e.g., in *Drosophila*, a Practical Approach (Roberts, ed. 1986)), and in Kernan et al. (1994), respectively.

Alternatively, it is possible to screen for molecules that can mimic *NOMPC* activity by performing the screen in a *nompC* mutant background. Those molecules that rescue the mutant phenotype can be considered potential MSCs.

#### Assays Using Oocytes or Cultured Cells in vitro

##### *Xenopus* oocytes

In preferred embodiments, MSC proteins are expressed in oocytes of the frog *Xenopus laevis*, and the mechanosensory transduction of the oocyte measured. Such assays are useful, e.g., to measure the activity of homologs, variants, derivatives, and fragments of MSC proteins, as well as to measure the effect of candidate modulators on the activity of MSC protein channels in the oocytes. In such embodiments, mRNA encoding the MSC protein, or candidate MSC protein, is typically microinjected into the oocyte where it is translated. The MSC protein, and in some cases the candidate MSC, then forms a functional mechanosensory transduction channel in the oocyte which can be studied using the methods described herein. In such embodiments, MSC cDNAs are typically subcloned into specialized transcription vectors in which the cDNA insert is flanked by *Xenopus* hemoglobin 5' and 3' untranslated regions. Transcripts are made from both the sense and antisense strand of the plasmid and then polyadenylated using standard techniques. These transcripts are then microinjected into *Xenopus laevis* oocytes. After allowing a sufficient time for translation, the oocytes are subjected to voltage-clamp recording. Cell-attached patches of oocyte membrane are assayed for the presence of conductances provoked by the application of mechanical force to the membrane, e.g., using small, calibrated pressure and vacuum steps applied through the patch pipette. Because *Xenopus* oocyte membranes contain an endogenous mechanically gated conductance, which is typically observed using these methods, the conductance due to the heterologous MSC channel represents any additional conductance, i.e., beyond the background level, seen during a mechanical stimulus. In such assays, it is important to compare the sense- to the antisense- and mock-injected controls for the presence of mechanically gated conductances.

##### Cultured Cells

In certain embodiments, MSC proteins are expressed in cultured cells, e.g., mammalian cells, and the mechanosensory transduction activity of the cell determined. In such assays, cDNAs encoding known or candidate MSC proteins

are typically subcloned into commercially available cell expression vectors, e.g., mammalian cell expression vectors, and then transfected into cultured cells. Expression vectors, transfection, and maintenance of animal cells are well known to those of skill and are taught, e.g., in Ausubel et al., supra, and Freshney, *The Culture of Animal Cells* (1993).

Cultured animal cells expressing MSC proteins, like the above-described oocytes, are subjected to cell-attached patch voltage-clamp recording during the application of mechanical stimuli such as small, calibrated pressure and suction stimuli to the patch. Osmotic membrane stress can also serve as a mechanical stimulus. Again, as eukaryotic cells generally contain endogenous mechanically gated ion channels, it is important to compare the transduction levels in the transfected cells to those in the mock-transfected controls. Any mechanically-gated conductance detectable above the level of the endogenous conductance is due to the candidate channel.

Alternatively, because MSC channels conduct calcium ions, transfected cells are loaded with a fluorescent  $\text{Ca}^{2+}$  indicator dye and then stimulated with hypo- and hyper-osmotic solutions while monitoring the cell's fluorescence. Hyper- and hypo-osmotic solutions create membrane stresses that open mechanically gated ion channels. In such assays, the influx of  $\text{Ca}^{2+}$  causes an increase in fluorescence of the  $\text{Ca}^{2+}$  indicator dye. As with the voltage-clamp recording, it is important to compare the transfected and mock-transfected controls. Any increased fluorescence in the transfected cells during the stimuli compared to that observed in mock transfected cells is due to the presence of the MSC channel.

#### Biophysical Properties of MSC Channels

The effect of a sequence alteration in an MSC channel, or of a candidate modulator on a channel, can also be assessed by examining the effect of the sequence alteration or the compound on one or more structural or biophysical properties that are typical of MSC channels. For example, MSC channels show very little voltage dependence, and are instead gated by mechanical stimuli. Further, MSC channels have a non-specific cationic preference, i.e., they conduct many different cations, including some large organic cations like tetramethyl ammonium ion (although weakly). The solution bathing these channels in the *Drosophila* bristle and in vertebrate hearing organs has a high potassium ion concentration (over 100 mM), which is very unusual for an extracellular fluid. Because of this, the principal current-carrying ion *in vivo* is  $\text{K}^+$ , with a small portion of the current carried by  $\text{Ca}^{2+}$ . In addition, as MSC channels are completely blocked *in vivo* by tetraethyl ammonium ions, it is expected that the channels are also refractory to tetraethyl ammonium ions in heterologous systems. Further, MSC proteins are in general refractory to  $\text{Gd}^{3+}$  ions, albeit at millimolar concentrations; in our bristle recording system, however, fly mechanoreceptor neurons are unaffected by  $\text{Gd}^{3+}$  treatment.

It will be appreciated that any of these characteristics, which are typical of mechanosensory transduction channels *in vivo*, can be assessed in cell-attached patches in either oocytes or cultured cells to assess the effect of any potential modulator, mutation, or treatment upon an MSC protein.

#### Candidate Modulators and MSC-Binding Compounds

Using the present methods, any protein, amino acid, nucleic acid, nucleotide, carbohydrate, lipid, or any other organic or inorganic molecule can be assessed for its ability to bind to or modulate the activity of an MSC polypeptide. Such candidate modulators or binding proteins can be deliberately designed, e.g., a putative dominant-negative form of an MSC

identified using high efficiency assays to rapidly screen a large number of potential compounds, e.g., from a library of nucleic acids or a combinatorial peptide or chemical library.

#### Proteins

Any of a number of polypeptides can be used in the present assays to determine their ability to bind to or modulate mechanosensory transduction activity in an MSC-protein expressing cell. Such polypeptides can represent, e.g., a candidate protein or collection of proteins encoded by a library of nucleic acids, can represent a putative dominant negative form or other variant of an MSC polypeptide, can represent a collection of peptide sequences, e.g., from a combinatorial peptide library, or can be predicted using a computer-based structural analysis program.

#### Heterologous Proteins

Polypeptide modulators of MSC proteins can be identified using a fluorescence-based screening strategy. In such approaches, cells are first induced to stably express an MSC protein, and then transfected with a cDNA clone of interest, e.g., representing a deliberately-selected candidate modulator or a collection of random clones such as a cDNA library isolated from a sensory tissue. The transfected cells are then loaded with fluorescent  $\text{Ca}^{2+}$ -indicator dyes and subjected to an osmotic stimulus or a mild mechanical treatment. Heterologous proteins that exert a modulatory effect on the MSC channel will cause the cell to exhibit either an increase or a decrease in the fluorescence during the stimulus compared to a cell expressing the MSC protein alone.

#### MSC Protein Fragments. e.g. Dominant Negative Forms

Because MSCs are thought to be part of a multi-protein complex *in vivo*, it is expected that a dominant-negative form of MSC can be produced by designing an MSC that lacks mechanosensory transduction activity but which can nevertheless interact *in vivo* with other molecules involved in mechanosensory transduction. A "dominant-negative" MSC refers to any MSC whose presence reduces mechanosensory activity *in vivo*, even in the presence of fully functional MSC protein. For example, overexpression of the ankyrin repeats alone (which are thought to facilitate protein-protein interactions), or in combination with a defective channel domain, will likely lead to the disruption of mechanical signaling. Alternatively, if these channels are comprised of several homomeric subunits (e.g., single MSC polypeptide units), expression of the channel moiety alone will reduce mechanosensory signaling in a dominant fashion.

In addition, because MSCs are weakly similar at a structural level to many voltage-activated channels, they could potentially contain an endogenous "ball and chain" inactivator of the channel (see, e.g., Antz et al., *Nat Struct Biol* 6(2):146-50 (1999)). Accordingly, one can potentially identify such endogenous modulators by producing small fragments of MSC, e.g., using a bacterial expression system, and assaying their ability to inhibit MSC protein activity in an assay as discussed supra.

#### Small Molecules

In numerous embodiments of this invention, test compounds will be small chemical molecules or peptides. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds that can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St.

Louis, Mo.), Aldrich (St. Louis, Mo.), Sigma-Aldrich (St. Louis, Mo.), Fluka Chemika-Biochemica Analytika (Buchs Switzerland) and the like.

#### Combinatorial Libraries

In one preferred embodiment, high throughput screening methods involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such "combinatorial chemical libraries" or "ligand libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Pat. No. 5,010,175, Furka, *Int. J. Pept. Prot. Res.* 37:487-493 (1991) and Houghton et al., *Nature* 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (e.g., PCT Publication No. WO 91/19735), encoded peptides (e.g., PCT Publication WO 93/20242), random bio-oligomers (e.g., PCT Publication No. WO 92/00091), benzodiazepines (e.g., U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs et al., *Proc. Nat. Acad. Sci. USA* 90:6909-6913 (1993)), vinyllogous polypeptides (Hagihara et al., *J. Amer. Chem. Soc.* 114:6568 (1992)), nonpeptidic peptidomimetics with glucose scaffolding (Hirschmann et al., *J. Amer. Chem. Soc.* 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen et al., *J. Amer. Chem. Soc.* 116:2661 (1994)), oligocarbamates (Cho et al., *Science* 261:1303 (1993)), and/or peptidyl phosphonates (Campbell et al., *J. Org. Chem.* 59:658 (1994)), nucleic acid libraries (see Ausubel, Berger and Sambrook, all supra), peptide nucleic acid libraries (see, e.g., U.S. Pat. No. 5,539,083), antibody libraries (see, e.g., Vaughn et al., *Nature Biotechnology*, 14(3):309-314 (1996) and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al., *Science*, 274:1520-1522 (1996) and U.S. Pat. No. 5,593,853), small organic molecule libraries (see, e.g., benzodiazepines, Baum C&EN, Jan 18, page 33 (1993); isoprenoids, U.S. Pat. No. 5,569,588; thiazolidinones and metathiazanones, U.S. Pat. No. 5,549,974; pyrrolidines, U.S. Pat. Nos. 5,525,735 and 5,519,134; morpholino compounds, U.S. Pat. No. 5,506,337; benzodiazepines, U.S. Pat. No. 5,288,514, and the like).

Devices for the preparation of combinatorial libraries are commercially available (see, e.g., 357 MPS, 390 MPS, Advanced Chem Tech, Louisville Ky., Symphony, Rainin, Woburn, Mass., 433A Applied Biosystems, Foster City, Calif., 9050 Plus, Millipore, Bedford, Mass.). In addition, numerous combinatorial libraries are themselves commercially available (see, e.g., ComGenex, Princeton, N.J., Asinex, Moscow, Ru, Tripos, Inc., St. Louis, Mo., ChemStar,

Ltd, Moscow, RU, 3D Pharmaceuticals, Exton, Pa., Martek Biosciences, Columbia, Md., etc.).

#### High Throughput Screening

In one embodiment, the invention provides solid phase based in vitro assays in a high throughput format, where the cell, cell membrane, or tissue comprising the MSC protein is attached to a solid phase substrate. In the high throughput assays of the invention, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) modulators. If 1536 well plates are used, then a single plate can easily assay from about 100- about 1500 different compounds. It is possible to assay several different plates per day; assay screens for up to about 6,000-20,000 different compounds is possible using the integrated systems of the invention. More recently, microfluidic approaches to reagent manipulation have been developed.

#### Computer-Based Assays

Yet another assay for compounds that modulate MSC activity involves computer assisted drug design, in which a computer system is used to generate a three-dimensional structure of MSC proteins based on the structural information encoded by the amino acid sequence. The input amino acid sequence interacts directly and actively with a pre-established algorithm in a computer program to yield secondary, tertiary, and quaternary structural models of the protein. The models of the protein structure are then examined to identify regions of the structure that have the ability to bind heterologous molecules. These regions are then used to identify molecules that bind to the protein.

The three-dimensional structural model of the protein is generated by entering protein amino acid sequences of at least 10 amino acid residues or corresponding nucleic acid sequences encoding a MSC polypeptide into the computer system. For example, the amino acid sequence of the polypeptide is selected from the group consisting of SEQ ID NOS:2, 4, and 6, and conservatively modified versions thereof. The amino acid sequence represents the primary sequence or subsequence of the protein, which encodes the structural information of the protein. At least 10 residues of the amino acid sequence (or a nucleotide sequence encoding 10 amino acids) are entered into the computer system from computer keyboards, computer readable substrates that include, but are not limited to, electronic storage media (e.g., magnetic diskettes, tapes, cartridges, and chips), optical media (e.g., CD-ROM), information distributed by internet sites, and by RAM. The three-dimensional structural model of the protein is then generated by the interaction of the amino acid sequence and the computer system, using software known to those of skill in the art.

The amino acid sequence represents a primary structure that encodes the information necessary to form the secondary, tertiary and quaternary structure of the protein of interest. The software looks at certain parameters encoded by the primary sequence to generate the structural model. These parameters are referred to as "energy terms," and primarily include electrostatic potentials, hydrophobic potentials, solvent accessible surfaces, and hydrogen bonding. Secondary energy terms include van der Waals potentials. Biological molecules form the structures that minimize the energy terms in a cumulative fashion. The computer program is therefore using these terms encoded by the primary structure or amino acid sequence to create the secondary structural model.

The tertiary structure of the protein encoded by the secondary structure is then formed on the basis of the energy terms of the secondary structure. The user at this point can enter additional variables such as whether the protein is membrane bound or soluble, its location in the body, and its cellular location, e.g., cytoplasmic, surface, or nuclear. These variables along with the energy terms of the secondary structure are used to form the model of the tertiary structure. In modeling the tertiary structure, the computer program matches hydrophobic faces of secondary structure with like, and hydrophilic faces of secondary structure with like.

Once the structure has been generated, potential binding regions are identified by the computer system. Three-dimensional structures for potential binding molecules are generated by entering amino acid or nucleotide sequences or chemical formulas of compounds, as described above. The three-dimensional structure of the potential binding molecule is then compared to that of the MSC protein to identify molecules that bind to MSC. Binding affinity between the protein and binding molecule is determined using energy terms to determine which molecules have an enhanced probability of binding to the protein.

Computer systems are also used to screen for mutations, polymorphic variants, alleles and interspecies homologs of MSC genes. Such mutations can be associated with disease states or genetic traits. As described above, GeneChip™ and related technology can also be used to screen for mutations, polymorphic variants, alleles and interspecies homologs. Once the variants are identified, diagnostic assays can be used to identify patients having such mutated genes. Identification of the mutated MSC protein encoding genes involves receiving input of a first nucleic acid or amino acid sequence encoding MSC proteins, e.g., a sequence selected from the group consisting of SEQ ID NOS:1-9, and conservatively modified versions thereof. The sequence is entered into the computer system as described above. The first nucleic acid or amino acid sequence is then compared to a second nucleic acid or amino acid sequence that has substantial identity to the first sequence. The second sequence is entered into the computer system in the manner described above. Once the first and second sequences are compared, nucleotide or amino acid differences between the sequences are identified. Such sequences can represent allelic differences in MSC protein encoding genes, and mutations associated with disease states and genetic traits.

#### MSC Genotyping

The present invention also provides methods to genotype an animal, including a human, for an MSC gene or protein. Typically, such genotyping involves a determination of the particular sequence, allele, or isoform of an MSC gene or protein, using any standard technique as described herein, including DNA sequencing, amplification-based, restriction enzyme-based, electrophoretic and hybridization based assays to detect variations in genomic DNA or mRNA, or immunoassays and electrophoretic assays to detect protein variations. The detection of particular alleles, sequence variations, isoforms, etc., is useful for many applications, including for forensic, paternity, epidemiological, or other investigations.

In addition, the detection of certain alleles or protein forms is useful for the detection of a mutation in an MSC gene in an animal, and is thus useful for the diagnosis of mechanosensory transduction channel defects in the animal. Such mechanosensory defects may underlie any of a large variety of conditions in animals, including conditions associated with impaired hearing, touch sensitivity, proprioception, balance, and other processes. In addition, mechanosensory defects may be associated with a loss of contact-inhibition in cells, and thus may be associated with cancer in the animal.

In particular, it has been discovered that mutations that introduce a premature stop codon into an MSC gene within the ankyrin repeat region, or mutations that remove or substitute a conserved cysteine residue between transmembrane segments 4 and 5 of the protein, result in a dramatic decrease in MSC activity and are thus useful markers for such analyses.

#### Pharmaceutical Compositions and Administration

Mechanosensory transduction modulators can be administered directly to the mammalian subject for modulation of mechanosensation in vivo. Administration is by any of the routes normally used for introducing a modulator compound into ultimate contact with the tissue to be treated, such as the inner ear or other mechanosensory tissue. The mechanosensory modulators are administered in any suitable manner, preferably with pharmaceutically acceptable carriers. Suitable methods of administering such modulators are available and well known to those of skill in the art.

Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions of the present invention (see, e.g., *Remington's Pharmaceutical Sciences*, 17<sup>th</sup> ed. 1985)

#### Kits

MSC proteins and their homologs are useful tools for identifying mechanosensory cells, for forensics and paternity determinations, for examining mechanosensory transduction, and for diagnosing mechanosensory defects in animals. MSC specific reagents that specifically hybridize to MSC protein-encoding nucleic acid, such as MSC specific probes and primers, and MSC specific reagents that specifically bind to the MSC protein, e.g., MSC specific antibodies are used to examine mechanosensory cell expression and mechanosensory transduction regulation.

Nucleic acid assays for the presence of MSC encoding DNA and RNA in a sample include numerous techniques are known to those skilled in the art, such as Southern analysis, northern analysis, dot blots, RNase protection, S1 analysis, amplification techniques such as PCR and LCR, and in situ hybridization. In in situ hybridization, for example, the target nucleic acid is liberated from its cellular surroundings in such a way as to be available for hybridization within the cell while preserving the cellular morphology for subsequent interpretation and analysis. The following articles provide an overview of the art of in situ hybridization: Singer et al., *Biotechniques* 4:230-250 (1986); Haase et al., *Methods in Virology*, vol. VII, pp. 189-226 (1984); and *Nucleic Acid Hybridization: A Practical Approach* (Hames et al., eds. 1987). In addition, MSC protein can be detected with the various immunoassay techniques described above. The test sample is typically compared to both a positive control (e.g., a sample expressing recombinant MSC protein) and a negative control.

The present invention also provides for kits for screening for modulators of MSC proteins. Such kits can be prepared from readily available materials and reagents. For example, such kits can comprise any one or more of the following materials: MSC protein, reaction tubes, and instructions for testing MSC activity. Preferably, the kit contains biologically active MSC protein. A wide variety of kits and components can be prepared according to the present invention, depending upon the intended user of the kit and the particular needs of the user.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of



clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

## EXAMPLES

### Example I

#### Chromosome Mapping and Positional Cloning of MSC Genomic Region

To identify mutations with potential roles in mechanosensory transduction, a genetic screen was carried out to identify mutations in *Drosophila melanogaster* that result in uncoordination phenotypes. This screen yielded mutations in numerous genes. Further characterization of these mutations using electrophysiological methods determined that several of the genes also reduced or eliminated bristle mechanoreceptor potentials (Kernan et al., *Neuron* 12:1195-1206 (1994)). One of these mutations, responsible for the *nompC* (for no-mechanoreceptor potential), present on the second chromosome, abolished nearly all of the mechanoelectrical transduction in mutant cells. Flies with this mutation are uncoordinated to the point of lethality. Based on these phenotypes, the gene underlying the *nompC* mutant was identified as potentially encoding a protein playing a central role in mechanosensory transduction, such as a mechanosensory transduction channel.

To determine the position of the *nompC* gene on the second chromosome, *nompC* mutations were genetically combined with various second chromosomal deletions, and the resulting transheterozygous flies were screened for lethality. In this way, the chromosomal position of the *nompC* mutation was mapped to a small interval on the left arm of the second chromosome, corresponding to map positions 25D6-7.

To physically isolate DNA in the 25D6-7 region, the proximal-most clone from a chromosomal walk in the nearby 25D1-4 region (George & Terracol, *Genetics* 146:1345-1363 (1997)) was used to probe a *Drosophila* cosmid library (Tamkun et al., 1992). Overlapping clones were used to "walk" to the area that contained the *nompC* (MSC) protein encoding gene, by mapping the cosmid clones to genetic breakpoints. At the same time, the cosmids were tested for the ability to rescue the *nompC* mutant phenotype. One cosmid was found to rescue the lethality, uncoordinated behavior, and physiological defect of the *nompC* mutation. This cosmid was thus determined to likely contain the MSC protein-encoding gene.

### Example II

#### Sequencing of the Rescuing Cosmid and MSC Gene

To determine the sequence of the cosmid containing the MSC protein encoding gene, the genomic DNA insert from the cosmid was isolated, sonicated, polished, size-selected, and the resulting 0.7-2 kb fragments subcloned into plasmid

vectors. Plasmids were purified and analyzed for the presence and size of inserts, and 123 clones with inserts of greater than 0.7 kb were sequenced. The sequences determined from these inserts were used to assemble large contiguous fragments, which were extended by designing ad hoc primers from the ends of the fragments and using the primers to read additional sequence from the cosmid DNA. In this way, the entire 33.6 kb cosmid insert was sequenced.

The MSC protein-encoding gene was identified and characterized within this 33.6 kb cosmid sequence using exon analysis, BLAST searches, and secondary-structure prediction programs. These analyses established that the MSC gene is a large gene comprised of 19 exons, encoding a protein containing at least 21 ankyrin repeats and a set of as many as 11 transmembrane domains (6 of which show significant robustness), that is weakly related to the TRP family of epithelial cation channels (see, for example, Montell, *Curr Opin Neurobiol* 8:389-97 (1988)).

### Example III

#### Sequencing of *nompC* Mutants

To assess the molecular defects of the *nompC* mutants, we used PCR to amplify the genomic DNA encompassing the *nompC* locus from flies with one of four mutant *nompC* alleles. In this way, all four alleles of the *nompC* gene were amplified in approximately 2 kb fragments that covered the gene interval. These fragments were then sequenced. All four of the *nompC* alleles showed mutations in the coding region when compared to the sequence of the cosmid and to the parental, wild type DNA.

In three of these alleles, the *nompC* (MSC) polypeptide encoded by the mutant gene was prematurely truncated in the ankyrin repeats by the introduction of stop codons. The fourth allele had a missense mutation between transmembrane segments four and five, resulting in a C to Y substitution.

### Example IV

#### Identifying MSC-Related Genes in Other Organisms

To identify potential MSC-related genes in other organisms, we performed sequence comparisons between *Drosophila* MSC sequences and nucleotide and/or amino acid sequences present in various public databases. In this way, a previously unknown *C. elegans* genomic sequence was identified as an MSC homolog. This genomic fragment was found in the "unfinished/orphan" domain of the *C. elegans* genome project database. Using a variety of sequence analysis programs, putative coding exons, intron sequences, candidate transmembrane domains, and homology regions with *Drosophila* MSC were identified. FIG. 1 shows an alignment between the *Drosophila melanogaster* and *C. elegans* MSC homologs.

Three signature sequences for MSC, based on alignment analysis between the *Drosophila* and *C. elegans* sequences, were identified and are shown as SEQ ID NOS: 7, 8, and 9.

---

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1  
<211> LENGTH: 24358  
<212> TYPE: DNA



-continued

---

```

<213> ORGANISM: Drosophila melanogaster
<220> FEATURE:
<223> OTHER INFORMATION: genomic nompC (no-mechanoreceptor potential C)
nucleotide sequence

<400> SEQUENCE: 1

gtgaccatgt tgcgggggac atgtttagta attgcaaaat cgatcaggtc tgggattttt    60
cttgggtctg ctggccagta tgtaggettta cccggggata attcgctctc ttaatgtgat    120
aatattaatc tcagaataat gaaaatgtca ttggtgtggg aaaatgtggg aaattgtcaa    180
ggaacgtaga gagtaacatg gtaattctat attttatttt tatttttctg atggtaaaaa    240
agttctagct ttatagtaat aatatcatta ccttgagtta gtaagattta aaaaataaaa    300
taagctgcat tttaaaagcc accttactg gttagacgac agcaacgata agataagttt    360
acatTTTTgc tacttgcacg acttgttgcg gcatcactga taagcaaaaca gacataattc    420
gcgtggctgg aggttttctt gattcctatc gctatatttc tgctcttctc atgcccccaa    480
aaaagtctg cccatactca aagaattgct ttttatttag ttgacctgtg tgcacaaatca    540
gcaaggcata tttatatctg caattggaac tacaattgat gcataagaaa tgagggtgtt    600
gtgaatatct ttgaaactga aacgaaagt agtaacttag tttagtaact agtttgttta    660
gatataagtg agttataagt tgaattaaa gaaggatcac ttcttctagt attgataaaa    720
ccatttatta tacagagagt tatagaagtg gctccatgta acctagacta gccaaaaaac    780
tattaggcat tcatttctct ggccacttgg gattttcgcg accagtcagc aaggatgaca    840
tactcccaat tgcgtctggt gcccatttgg gtttccacc ggcaactaac gacgttgga    900
atcccaacga aacttaagag tagcgtccag attttggcgc caaaaaaggc ggtattattc    960
ggattcaaca attgtaaaca aacgcttgcg cggatgccac ttggctctta cctctgattt   1020
ttcgcaggag cgtcttgggt ccttcgagtt tggagcttcg tcgtgttgcc agagctacca   1080
aaccgagtgg agggccgagt ttttcgctc gagcgccttg ggaatagtcg actctgtgaa   1140
aatgggactg gcaaatcaga aactcgcaga cgctcgtggc aaacggttga ttttttctc   1200
gtcgtccga aaaaaggcaa aatagtaggc aacctgaaat ccagagtgtg agttggggac   1260
tcttttggcc aaaaatacaag gaggagaaaa atagaaaata ataaaggggg caccgccgtt   1320
aacgcacacg caaccgaagc cataaagggg ctaaacatat aaatttgtgt agtaaaagtg   1380
aagaaagcga aagaatcaaa gtggaataat agcgagtgtt tttcggtttg ctagtgtgtt   1440
tctgagtcgg agtttgtgtg tgtgtgtttg tgtgattcct agtgtgtctg ttgctgttgc   1500
caatgaaat gcaattgtt ggtaacaaat attggtaaaa tgcggaggcc gtaggaattt   1560
gtgcaatgcg agtgcaagt gaaggagccc gaaactatgc agctaaaaac ccgccatcct   1620
acccgcacg gaatcaataa taatacaata acccaaacgt attacacgga taatggcagc   1680
ataaaccagt taacatccga cagtgtttcc gactaacctc agagcaccta gctcatcccc   1740
cctgccacca accctcgaa aatcccat gatcagcgc ggattgtgga gcagtaacta   1800
gcgaggcata ccaggatgct gcagcgcgc ggaggcgtg gcggtggcg tggcgcgga   1860
gtgggtcgca aaaccctc ctcgtgacc ggcccaccg atgagtcggc tacgccagc   1920
gaacgggcta cgcccagc caaagcagac tccgatcca aggacgatag ctcgagcaat   1980
ggcgacaaga aggatatgga tcttttcca gcccacaaagc cgccgagtgc cggcgctcc   2040
attcgggaca cggcgaacaa ggtgctcgga ttggccatga aaagcgagtg gacgccatc   2100
gaggcggagc tcaagaagct ggaaaagtat gtggccaatg tggcgagga tggcaatcac   2160

```

-continued

---

atacgcgtgg	ccggcgttca	cgacatggtg	agtactgtac	agtgaagtgc	cgcgaggcgg	2220
gctttccggc	tcatttgcct	cgttttgtaa	aatcaattgc	gagccaaagc	gggaatagga	2280
agcgaataaa	atacaggaac	aggccaaca	ctcagcgaac	aatatggtaa	attaaatgta	2340
tacctagaga	aggattatca	atagttttaa	taaggttatt	gaaatcttta	aaactataat	2400
ttctatggat	cttttagttg	tatttatttg	aaaaatttcc	ttaagttttt	gtgtaatatt	2460
tcctcgagtg	tatgcatgtg	agaaacgtcg	cccttatcaa	cgtcggcggc	attttcccat	2520
ttctggttgt	ttaccagcca	aaataacgac	acaggaactg	gaggccagaa	aacagagcac	2580
accatggttt	ggccaaaaa	cagaggctag	caaggaaaag	cgcccaaaaa	aaaaaaaaac	2640
agagaacagc	gaatgttatt	tgatagctcg	gcccaaatgt	tttggctgcc	aaggcgatgg	2700
ctttggtggc	attcggtttt	gtagctccaa	gttcctgaag	cgctctgcca	caagttgcgc	2760
cgatacgcct	ttggggttag	cccccgctc	gaccgataaa	ctcataaaac	atcgaagaat	2820
tgaagcgctt	cgatttcaat	ttaccataaa	cgctatgaaa	cggagaagtc	gttgacataa	2880
aattaacggt	gcaccgctaa	tgaatgctgg	ggagggtgtc	ggcgaaggg	ttgaaacttc	2940
ctggcagggg	ttttctttta	cttttttct	ttcctttttt	ttttgtgtg	gtactatata	3000
tcccaactag	atgtgcagg	tgtctgctag	actagactta	cgacgagacg	gtatttgcat	3060
aaatatagct	tggagttgag	ctatttttgc	cttgattatt	tccgctttcc	cagaacgggg	3120
gtctttatc	ggttcttgac	ttgatgggct	tgctcttgat	ttcgttttaa	ttacgagcca	3180
acgagcttat	aatatcacat	ccagcttatt	agccgaagga	ttctaagtca	ataaagatga	3240
atttaaatgg	ccaagtgtct	tttcaatgag	gtcagcgggt	tggaaaggaga	gtaccatgta	3300
ttggactat	gttattgtgt	ttaaaatgtg	catatattaa	tattgtatta	ttcttacctt	3360
aagcttaagt	aatccccata	catttccatt	gcagaatacc	ggcatgacgc	cgctgatgta	3420
cgcaacgaag	gacaataaga	cggccataat	ggatcgcatg	attgagctgg	gcgccgatgt	3480
gggagccccg	aataatgtga	gtcttgagcg	ggaatagggc	aggaaataat	taaagcacct	3540
tagccaacte	cccacgggtg	tgggtccaaa	tatagaagcg	gccagctgt	ttaagccaac	3600
ggcggcagca	aaagccgcta	aaaatgtgtc	aatcaataa	aaaccgata	attaaatctt	3660
gagcgggggc	gttgggtggg	aaactcgtgc	accacttct	acgcacgatt	ctcacacgcc	3720
gcccaccacg	gtcaaatctt	caattcggca	atactcctc	gccgcaatgg	gtcaacttgg	3780
caggacttgg	ccaatgggta	gttcgcttca	tttgactcca	gttgagtaa	gttttccagc	3840
acgaatggga	atctcctcaa	gaaaaagaaa	tactaacaca	ttgcttttat	tttcatttta	3900
taactgctaa	caaaaaatta	taactctta	tttatagaaa	actaaattat	tattgggcac	3960
ccctcgtttt	taagtggctt	aaagttcgaa	cttaactttg	gtttttaaag	aaacagcaag	4020
tattactcat	aataatgtaa	ctcaacaaaa	gagttttccc	aaagagtaga	gatgtaaggt	4080
catcgctgat	gactatcctg	atctccccc	taatttacca	tcgtgattat	ggccaattct	4140
tttttttttt	tgatgtcagc	aagtgaagtg	agccaggttg	gcacgcccc	ttaggccaag	4200
ttgctaacaa	ttggtcgaat	tcgcccagca	gcttgctttg	catgccgcaa	ttacttagca	4260
catttcat	gaagtcgctt	tcttggctgc	ccattcacat	gtccttacgt	atagcgaacg	4320
tactttat	cggtgctagc	ggcgataaaa	atccttgacc	taattacaaa	ataattgttg	4380
ccaaaccagt	gcagacatgg	cgaattgaa	tacaaaaaca	aacacagaaa	gttcaatttt	4440
cccttctccc	ttgaaaaatg	ttctcctaaa	agattaaaga	gtgtgtaggg	aaaatgttaa	4500
aggtaaattt	gcacatgaaa	gtcataaaac	attaactagc	cgggagttac	aagctaagca	4560

-continued

---

tgaaaataaa	acactcgata	agactttata	tgagtataag	aatttatfff	cgttttaaca	4620
ggacattcat	tacacaaatt	ttgccaatga	tacttgggtg	tttaaaatat	tgagaaaatg	4680
ttgtccaaac	tgcaactaaa	aaccacatat	atattaatta	attatattta	atataaactt	4740
tccctttttg	caacacaatt	aattatgata	attattcatt	ttaaaactgt	tccatttgga	4800
tgattgttcc	ctcttgttgt	tcagctaatt	aaatattatg	atatcatttt	cgtaggttta	4860
tacaaagcgc	acctttttga	aaaccattac	ctcatctgta	taattactct	ttgttttta	4920
taaaacaaat	gtcacttcgt	gaccaaatcg	gataatttcc	cttacctga	ccaatgaat	4980
taaaaactga	gaaatgttta	ttgcatttac	aattcgcaac	ttatctaact	gtcaggtctg	5040
gtccaaagta	ataccctaac	aacacgacag	gaccaggacc	tttatggcca	ttataaagga	5100
tactcgtatg	atgtaacgcc	gtggttaatta	acatttttaa	cttttcaact	gcaaggtggc	5160
agactgcttt	tttttcggca	ctcgacttgg	aggcgtgctc	gcaacacctc	tttgaacgt	5220
aaaagccaat	taatcaagca	catgactccg	atgtacgcc	agttggccaa	aaactccatt	5280
tgacctttcg	agtgtggccc	aaaccggaga	cctcgacgtc	ggccccgact	tccgctacat	5340
ttttatggcc	agcggcgtca	ttaatatgca	attttaatta	aattcaagtg	gaattcttca	5400
cgcagtgacc	cctgcatatg	tgtgtggcga	tgacagcgtg	aactaaaatg	aggaataaaa	5460
acgccaatc	atttgtcaag	ttgcctcagt	gctgtagtga	agtaactctg	cccatccaag	5520
caaaaaaaaa	gcaaatat	tcacttcatt	agaaagtgg	gcacatgcaa	gaaggtggag	5580
ggattaagcc	aatgagcac	cgtaatgagg	acttgcaatt	attccaaga	aggtgtgtga	5640
catcgccaga	aatgacttc	atggcttcca	cgcgactatc	cccagatc	tctgggccgt	5700
aaaaacaaaa	caccacgaa	actgggtcga	cttcgtacac	ccttatccac	ccaaccttat	5760
cccttttcca	tttggcagg	caaaaatgtg	ctggaaaatt	tgcgcttccg	ctttgtttt	5820
gtttccgggt	tttcccttcc	accagccaag	caaacgcaaa	cacaagcgca	caaacacaca	5880
agactcgaaa	acgaactcga	acctggctca	aaagtatgca	aaacagcgcg	tgaatatatta	5940
tctgtctacc	ttggaagcca	atgcaacccc	aaaccagcag	cgattccgcc	caccgcgcca	6000
agtggctgaa	agtttacttt	gctttttctt	tagggccaac	acgtcttggg	tgggctttct	6060
ggacatgtgt	caaagccgtc	gactccgagc	gccaacttgc	gttgtatgca	aattagcagc	6120
agctgcggcc	agaaatagtc	gcaataaac	cgcagggaac	tcgaatttca	cacggcacga	6180
agccacacaca	cactgactta	agtgggaaag	tttgaaatc	ccatttggat	tctaggaatt	6240
gtaaaaaatc	atgtgcaaga	acacatagaa	tgtataaata	tagaattatt	ttaaatggca	6300
taacttctgg	tattctccta	attttttaac	atataatcta	aactaagtat	tattttcctt	6360
tcactatfff	tattaactag	aaattogtat	ccttttatgt	tgaatfffgt	agactctgtc	6420
tgcaacttacc	aaactgatga	cagggccaaa	agcaccata	cataatgct	aaaccagttc	6480
acttccgttt	tccgggctaa	gaactgtggg	gaggttagt	tataaattag	agccatggtc	6540
cgaggtccga	gcatacgggg	cgtatgtgta	acacgttgcg	ttatggctta	ttatataagg	6600
caataaatat	ggccaaatgc	ccccgattca	tatgtgactc	acttggctat	tagctggcgt	6660
taaaactaagc	actccatgtc	agacgttatc	ttaaagcact	tttcggtacg	tttcggtgat	6720
ttgtccagg	tcatatfff	ctagccgat	tgttttatat	ttcttttcgg	gttttctctg	6780
tccgcatgta	tcagttfff	gcattgtgag	ttgcggctgg	gctgtggcca	ttaagaaaac	6840
cccgtccgta	agtgaagtc	cgcattgcaag	attgtggctt	aagtaatcaa	ccactccctt	6900

-continued

---

ttgccccgtt agcccgatgc aaaaccgact gactttgacc cattgaaactg acccagctct	6960
tttgggtgtgg gggcgctcagt ttcctgccaa tgaattgcaa ttgatttctt cgtttcttct	7020
cttctcttct ctttcaggat aattataatg tgctacatat tgccgcaatg tattecgcgtg	7080
aggatgtcgt caaattgttg ctaacaaaaac gcggcgtgga tcccttctcc accggtggcg	7140
tgagtattcc aatagcttta tatactacat atatacgtat gcgccccaaag aaagtgttac	7200
cccaatagtt gaggtagcga cacgtcaggc gacacactca atactcgagt tcctactttc	7260
gagtcaatga aatagctgca taccttgggg ctgctgtcag cccgattcgc aggcaatttg	7320
cggtcattag acgcatactt cacctggcct cgaaagagaa gaaaaaaaa aaacctatcc	7380
aaaggtcaga gccatgcgaa gatgcaactt tgaggctcgc atgttgcatg ttactttggc	7440
gggaccagca attaactggc gacaaggta agatggtaat gtctagggcc cgcttaagaa	7500
cactttaaga cctgaaaaca aatttaaagt aaccctaggt ttcacgaaaa actttactca	7560
tcagattaaa cagaaattta agcttagata ccgtcattaa aattaaatt taacattttg	7620
catgatttcc aagtctgact tctgtttaaa tactacaatg tataaatatt aaagtctgag	7680
caagattagt gacaccatct ttatattgtc taaaatcata aagcgttaac catttaatac	7740
aatgcatttt ctcataggta acatttttaa caaaatata gatgatcaca tegtcaagca	7800
ttttggcaat tttttctcca agtttatttc tegtgtcggc attaatgtgc ttttctttat	7860
ttttttctcg gcgcatttg gttttcgaga cttggttatt tagggggcgt gcgccttgc	7920
caaaactact atggttatca gaagagagct ctaagcacgt gtgggagcga gagaagtgga	7980
gctgcggaag cgagacagac agatgcaaac ttttgtttta gcaacagcca agtttgaagt	8040
gttccgttag cgtgtgtgcg tggcaaaaag gactcccaca tccacaaccg acacctgccc	8100
cccatgttgc ctacacctgc tgctcgacca cccctcccc accatcacct atatacact	8160
ctctcgctca ctcccgcagc ggttgtcggg gggagttctt tattatgctt ttttcgggct	8220
gtcaatctgt gatatgagcg ggagaggcca aaaaagaaaa atgacacgaa atgtgcttat	8280
aaacgcaaaa acgagccact tgcctattca gtagcaaatg gaattttgaa gcgaataggg	8340
aaacagtttg ccagtttttt aggtgccaac attaaccaaca cagtgtgca catagctgca	8400
tattaatttt ggctagaaaa aaagtgtaac cccagcaata agtgcgtttg cagtgtgtgc	8460
atagtttaat cgaagactta attggatttt tttccctttt cagtccgctt cgcaaaactgc	8520
ggtgcatttg gtgtccagtc gacaaaaccg aactgcaact aatatcctgc gcgctctgct	8580
cgcggcagct ggcaaggata ttcgcttgaa agcggacggc gtaagtgtta ccatgtgtgc	8640
ttgtgattga gtgtgccagt gtggctgtgt gtgtgacgag gagagccaca agtgttgccc	8700
gcccattga tgcgcttta tctccactag tttatgatag ctaagccacc caaatgcaag	8760
ccgagtgtga gtcaagtact ctgcacagcg gtgccaggcg gtgccagct aaacaaagac	8820
ttaataaaaa tcaccaaaaa atatatacat tacaataatg gcaccaacaa aatcgagagg	8880
agttagtaac ataaagcaaa caaaattgtg tggaaaaatc gatatgcaaa actgctcgcg	8940
gtaaatgcat ttcgactggc tgtaaatcag aaaaggccca aaaaagttaa tgcggctatt	9000
acacagcgag gaattgaata ggtaattttt gagtcaattt tagcttataa tttgtggtac	9060
ttttatgaat ttttttaaaa tttttatttc aaattattag agagctaata tatttgaatt	9120
atgcttatat aacttaaaat actcaaaatt tatagacagc aataaagtat gggatctgca	9180
acacatcttt ttctacactg tatcaataag tagctctcac cacagtgggt aggctccagc	9240
gagctttgaa ttaccatcga agcagttgtc tccgctgat gaacttctg gggctaaccg	9300

-continued

---

agctccagat	ccctttttcg	agctccccc	ttgaaatct	gaacagaaat	gcggaactat	9360
ttgtgcate	acgtgcccc	ggtgaaaatg	cacaggggat	atttcatta	cgcacgcgaa	9420
gaaagcgcat	aaatttccaa	cgaattgcta	tcaagcgatt	gtaaggattt	ggggtatatg	9480
ggggctgatt	gagggaaatc	cggtgccc	cgattgattg	tctagacaaa	atgggtaacc	9540
cacctcgatt	tgtgcctcga	gggctgccc	aatggcaaa	cagcaacttg	atttaaatca	9600
attagagaga	ggtggaatgg	cactgtcagg	cgaaattagt	cggatgaagt	atttagcttt	9660
cgatggcatt	cagttcgatt	cgtttcgatt	cgcttttctt	tttttttct	acacgcattt	9720
ccggtgtgca	tatacatgca	aatatatata	ttgtatgtgt	gtggatagta	ctgtagtttt	9780
cccccgcgag	ggcgtcaac	tcggtgccaa	caacaacaa	atataacaaa	gcgaggaaaa	9840
ctctaccgaa	aaaaggggt	caagtcgctg	tacaacttga	tttactcgcc	tttctggca	9900
gatagggata	atggctccc	gtcacgccc	cctcttacga	ctcgcccca	aaaggtagtt	9960
ggttgcaagt	tggagcgcca	aagttgcgaa	cttggctaaa	aatagcgaaa	catggtgccc	10020
ttaacacttg	aggctcgaat	tggctaattg	gatatttatg	attatatggt	cgcgagtgtg	10080
aatggatgtg	tgttcgctgt	ccttatctta	attatatttt	atactatata	taacctatct	10140
ctaacctagc	gtggcaaaa	accattgtct	ctggccgtgg	agtcgggcaa	ccagtccatg	10200
tgcagggagc	tctgtgctgc	acaaacagca	gagcagctca	aggtaagtaa	tctgtgaact	10260
agcagataag	tttaccact	tattttaaaa	cctaaaagtc	tagttgcagc	ttatattgat	10320
ttaaatagaa	acactgaata	catcatctag	ttaataacca	aaaatgtcaa	cagtatgagc	10380
cattaaaagc	ataaaatgct	aatttcttat	accatctacg	catctaactg	atttcctaac	10440
taggaccaag	aaattgttga	ttttataatc	gccacgatag	tgtcaatcaa	actgtccatc	10500
tgagctgtcg	gaaaatgtcc	acaaggttct	taaagccttg	aactgtccaa	taaccaagcg	10560
tgtaaataaa	tcaaaaatgc	aaatttacc	tgctcacctg	tgcgtacagg	tgcattgcaa	10620
gtgcaacagt	gcgagcatt	ggcaaaagtt	gtgcaatttt	caatcagaag	ttgaagtgca	10680
acacaccaag	agcagtgctg	gttgattaaa	ttaaccaaa	ggctacggct	cgcttcaggc	10740
caagggttca	agcccaagtt	aaagttaaag	ttgcgcctga	ctttggccgc	tggctgagca	10800
cgcaatcagc	cgcaaaaaca	gccgtaaact	gggtcaaaac	tgaggcgaaa	acgcagctaa	10860
gatgggaagg	gaatctgatt	tgcataagccc	aaaataaaat	gtcgaaagtg	aaatgcagca	10920
acactaagga	aaaatttaag	taaattattt	aaaaatattt	aaacaatgaa	gctatgaagc	10980
tctagcaaa	ataccaat	agttagggaa	tatcattata	atttgtcaca	tagttaatta	11040
atttcaagca	taggagcaat	tatgactttg	caattatata	aaaacatttt	tgtgaagtgc	11100
accctttcat	gttaaatttt	ggatttattt	tttcgcaggc	aacgacggcc	aatggagaca	11160
cggccttgca	tttggccgcc	agacggcggg	acgtggacat	ggtccgcac	ctggttgatt	11220
acggaacgaa	tgtggacacg	cagaatgggg	agggccagac	gccacttcat	atcgcggccc	11280
ccgaaggcga	tgaggetcta	ctcaagtact	tctatggcgt	gcgcgcctca	gcgtccattg	11340
cggaacaatca	aggtgagtct	gtgggaatgt	ggagcaagga	aaagcatggt	gcaaatcgtg	11400
tttgaccttg	atataacaca	ataaaaatca	tgaatttttc	acttctcaat	agaagctagt	11460
gattataaag	tggaggata	aagtatatgt	ttgtggcgcc	cccggttgga	ccgagctcca	11520
gacatacga	tgtccgtctt	gatgattaaa	atttatatat	atatatatgt	aataccctat	11580
agatcgcact	ccgatgcact	tggccgccga	gaatgggcac	gcgcacgtca	tcgagatact	11640

-continued

---

ggccgacaag	ttcaaggcga	gcatcttcga	gcgcaccaag	gatggcagca	cgctgatgca	11700
cattgcgtca	ctcaacggtc	atgctgagtg	cgccacgatg	ctcttcaaga	agggcgtcta	11760
cctccatattg	cccaacaagg	atggagcccg	gagtattcac	accgccgccc	cctatggtea	11820
cacgggaatc	atcaacaccc	tgctacagaa	ggcgagaaa	gtggatgtga	ccaccaatgt	11880
aggtgggata	atgtattaag	ggataatcgt	attaattcca	cactctttgc	aggataacta	11940
tacagcactg	catatagccg	tggaaatcggc	taagcccgcc	gttgtggaaa	ccctgctggg	12000
atgtggagca	gatgtccatg	tccgtggcgg	aaaactacgt	gagaccccgc	tgacattgca	12060
ggcacgagtg	aaggatggag	atagggtgtg	cctcatgttg	ctgaagtcgg	gagccagtcc	12120
aaatttgacc	acggatgact	gtctgacccc	cgtgcatgtg	gcccctcgtc	atggcaatct	12180
ggccacgttg	atgcaactcc	tcgaggacga	aggagatccg	ctgtacaaat	cgaatgtgag	12240
tagattatta	gaatagaatg	ataaacgctt	gaattaaac	ttccatttta	tagactggag	12300
agacaccgct	gcacatggcc	tgctcgtgctt	gccacccgga	tattgtgctg	catctcatcg	12360
agacggtgaa	ggagaaacac	ggtccggata	aggccaccac	ctatataaac	tcggtaaacc	12420
aggacggcgc	cacggcgttg	cattacacct	gccaatcac	caaggaggag	gttaagattc	12480
ccgaatccga	caagcagatc	gttcggatgc	tcctcgaaaa	tggtgcggat	gtcacgttgc	12540
aaacgaaaa	tgcttgag	accgctttcc	actactgcgc	cgtggccggc	aacaatgatg	12600
tgctgatgga	gatgatctca	catatgaatc	ccacagacat	ccaaaaggcc	atgaaccggc	12660
aatcatcggg	gggctggact	ccactgctga	ttgcttgcca	tcgagggcac	atggagctgg	12720
tcaataatct	actggcgaat	cacgctcgag	tggatgtctt	cgataccgaa	ggacgatctg	12780
ccttgcaatt	ggctgctgag	cgaggatacc	tgcatgtgtg	tgatgccctg	ctgaccaata	12840
aggcttttat	taactccaag	tcccgcgtgg	gacgcactgc	actacatctg	gcagccatga	12900
atggatttac	gcatctgggt	aaattcctga	tcaaggatca	caatgcagtt	atcgatattc	12960
taacgttgag	aaagcaaacg	ccgctccatt	tggcggcagc	cagcgggcag	atggaagtct	13020
gtcagctgct	cctcgagctg	ggcgccaata	tcgatgcgac	ggacgatctg	ggccagaagc	13080
caatccacgt	cgccgccag	aacaactact	ctgaagtggc	caaactcttc	ctgcagcagc	13140
atccatccct	ggtgaatgcc	accagcaagg	atggaaacac	atgtgccac	attgccgcca	13200
tgacgggatc	cgtcaagggt	atcgaggagc	tgatgaagtt	cgatcgatcg	ggtgtgattt	13260
cgccgcccga	taaacttacg	gatgccacgc	cccttcagct	ggccgcccag	ggcggacatg	13320
cggatgtggt	gaaggctctt	gtgagagctg	gtgcctcctg	caccgaagag	aacaaggcgg	13380
gattcaccgc	cgttcacatg	gcccacagaa	atggacatgg	tcaggctctg	gatgtgctga	13440
aaagcacaaa	ctcactaagg	atcaatagca	aaaagttggg	tctgacgccg	cttcatgtgg	13500
ctgcctatta	cggacaggcg	gataccgtgc	gggaattgct	gaccagtgtt	cccgccaccg	13560
tcaagtccga	aactccaacg	ggacaaagt	tatttgggga	tctgggcacg	gagtcgggaa	13620
tgacaccact	acacttggcg	gccttttccg	gcaacgagaa	cgtggtgcga	ctgctcctca	13680
actctgcggg	tgctcaagtg	gatgcggcga	ccatcgagaa	cgtaagatta	cctgcatatc	13740
tcttctgttc	agaaaccatt	aacacaacaa	ttgattctac	agggtataa	tccactccat	13800
ttggcttget	tcggtgtgca	catgtcagtg	gtcggtttgc	tcctaagtgc	gtcggcggaa	13860
ctctccaat	cgaggatcgc	taacggcagg	acgggcctgc	atcgcgccg	catgcatggc	13920
catatccaga	tgggtgggat	tctgctcggc	cagggcggcg	agatcaacgc	aaccgatcgg	13980
aacggttggg	cgccactgca	ttgtgctgcc	aaagctggcc	acttgagggt	ggtgaagtgg	14040

-continued

---

ctgtgcgagg cgggtgcctc gccaaaatcg gagaccaact acggttgccg cgccatttgg 14100  
ttcgcgcct cccgagggaca caacgaggtc ctgcggtatc tgatgaacaa ggagcacgac 14160  
acctacggcc tgatggagga caagcgatc gtgtacaacc tgatggtggt gtccaagaac 14220  
cacaacaaca agcccattca ggagtttgc ctggtatcac cagcaccctg ggatacagcc 14280  
gccaaaactgt ccaacatcta catagtactc tcgacaaaagg tgatttagct aaaggatctc 14340  
tatgcactta actaaactaa ctaactaaaa cttttgatc tctttaggaa aaagagcgcg 14400  
ccaaggatct ggtagcagct ggcaaacagt gcgaggcaat ggccacggag ctcttggccc 14460  
tggcagctgg gtcagattcc gccggaaaaga tccttcaagc caccgataag cgaaacgtgg 14520  
agtttctcga cgttctcatt gaaaatgagc agaaggaagt gattgcccac acggtagttc 14580  
agegatactt gcaagtgtgt gatattattg actagcttag atcttaactt attgagattc 14640  
tgatatgtat cttctctct acttttagga actctggcat ggctccctga cgtgggcatc 14700  
ctggaaaatc cttctgctgc tcgtggcctt catagtctgc ccaccagtgt ggattggatt 14760  
cacattcccg atgggtcaca agttcaacaa ggtgcccac atcaagtca tgcgtacct 14820  
aacctctcac atttacctca tgatccacct gagcatcgtg ggcataacgc ccatttacc 14880  
agtgtccga ttgagtttg tgcctactg gtaacgagtg ggtcttctca tctggctgag 14940  
tggattgctc cttttcgagc tgacgaatcc gtcagataaa tcgggactgg gatcgataaa 15000  
ggtgctcgtg ctgctgctcg gcatggccgg agtgggtgct catgtctcag ctttctatt 15060  
cgtctccaag gagtactggc caactttggt gtattgtcga aatcagtgct tcgcttggc 15120  
cttctgctg gctgtgtgct agatcctcga cttttgtcc ttcaccacc tattcggtec 15180  
ctgggccatc atcattgggg atctgctgaa ggatctggct cggtttttg cctcctggc 15240  
catctttgtg tttggctttt ccatgcacat tgtggccctg aatcagagct ttgccaattt 15300  
ctcaccggag gatctgcgca gcttcgagaa gaagaaccga aatagaggct acttcagtga 15360  
cggtaagtgc aaacgtttgc tttgctttct ccagtctact tttcgaattt ttgtttcgaa 15420  
ctttttgttt tcatttgaa tgttttgca aacttcctct tttgaacgtt caatgtgtct 15480  
tgataagtat ctgtgtctgc cttgaatgaa aagcccctct aatcaatgtg cgtcgtatgt 15540  
ttcacataag taaaataaag caaaaagaa ccaacttcaa ccacataata caacaattgc 15600  
atgctcaaca agtacaacaa acccgaacct ccaacctga tgcgtaatc cccgtccacc 15660  
ctccaccaaa aagacctcca ctaataatgt tctcctctg atcttaacct ccaactgaat 15720  
atcttaactg aattatccga atggaacaga tgacatgcc acaccccgac ctccgccggt 15780  
ggagaattat gtcgatagtc gcttcagcga attccgacga aagcacaagg acgaccgtaa 15840  
gtctcctacc atccacaact accaaccctt actaccccg catttgcatg gcccccctt 15900  
ccgggggctg ccccgcccc ttaacccaac aatgccgaa tccaaaccgt tgcgttgccg 15960  
ccttcgatgt tgtgcgtaaa gtgttaatgt cgtttgttt ctagtccct ggaggaacat 16020  
ccacaagtcc gactcgtg ctcgaaatcc cctcgccttg ctagtctcag ttaacttctg 16080  
ttgaggcat gttcgcggga aatcccttt tccgcatcct cgatgttggt gatctgtgtt 16140  
tatataggta tccatgcgcc aagctttatt acttagtttg gtagatcgtt ttatacctt 16200  
gcttgatca atttttaatt atatgtatct ctttatgtat ttttaagtga catataaata 16260  
caaaaaaatt attaagaatc agaatttaaa accataattt attctcatta aattcaatca 16320  
ttattatttc aaaaaatcct agatctgtgt ccgatattat tttcttact atatttgta 16380

-continued

---

```

ttctttttt aagtttagatt ttttatcgat gtgtaaccag agcgatatcc attagaactc 16440
tgtacaaact aaaaattcca gtaatgcatg ttgatgtttt tatccagtca atccaaacca 16500
aaatcaaaaca atcaatcagc aatatcgata taaccaatgc cgcctgcct ggggctttca 16560
gcttgccgag cttgcccacc accaaattct gcacaatcga aacaatcgag accgatcgaa 16620
tcgaatcgat aacgaaaaac gataacgcta ctgataccga ttaccgatgc tcgtattcgt 16680
gagtcattcg aaccgctcag ctgcgaactg cgagatgctg cttttgacgt gtttaaccac 16740
tcacccgcac tctccaaaat ccaaataaac ccaccataa atatactcgt ttaatgtaaac 16800
ttcaaaataa ccaacaaata ccaagtatta aactcgaca cacgcctgtg ccaagccgac 16860
aatatatata cgtatatata cgctagctgc agcaatcgca atgcaatagt tcagttatct 16920
gattgtgagt aacgttccgt tcggaccat gttaggacc atgacgcct tctggcttt 16980
cgagcgctc ttcttcgagg tcttcggaca gacgaccacc ctggacatca atccatgag 17040
acacttgctg cccgagtgga cggaggtgct cttcaaattt gtctttggca tctacttgg 17100
gggtgctgtg gttgtactca ttaacctgct aattgccatg atgtcggata cttatcagcg 17160
cattcaggtt tgtattgcca aggccactaa tcagtatttt ctctctgctt tccctcttc 17220
ccggtttatt tgtttcaatt ttcatttacc ggaatgctat ttgtttgtgc tttgattgta 17280
acaaccccaa aactgaccgc tccaaattga aacacaattg ggcatagaacc gaaactgggg 17340
gttggctgat cggacaaatc aacgaaacaa aaaaaaaaa aaaaaccaca taatcgaatc 17400
aaaccaacca acctggggct cggttatctt tttattttc aaaataattt ccacgcgggc 17460
caatatatgc gtgctgtccg ggggtgtcta tttgtatctg tatctgtatc tggaaatgta 17520
tctatgggtc tccgacacag tgcgcatgca tccgattaac tcgttcgagt tgttgttctt 17580
cgcggtgctc ggacaaacga cgaccgagca aacgcaagtt gacaaaatca aaaatgtagc 17640
cacgcccact caaccgtatt ggggtgagta cctgttcaaa attgtctttg gcatttacct 17700
gttgggtgctg gtggttgtgc tcattaacct gctgattgct atgatgtcag acacctatca 17760
acgcattcag gtagtattgc taaatgcgct tttatctaac tcgactctat ttattaactc 17820
gtactttaac cataagtata taaatttcat attgcattgt gtattaatca ttctctatct 17880
cagcataaga agtaaattha catatgaaga tgatttatat ttcttagata tataatagcg 17940
gtagttagga agtgagctgt tttgggaaca tattgagaaa atagtttaatt aatctggaga 18000
acttggcatg ctctgtaaat ccatcaactg ccagacttg catcttcag gttttttcag 18060
gaaaataatg ttagcaatct gagggataca atttgtgaa agtgtatctc aaagatggaa 18120
gctgcccgc ttctagtgtg gtacagtgca gagtagcttt agtggattag cgccttgaa 18180
gtgtgccctg cttttgtgac cagtgttgag cgaggccaaa ccagaaagtg ttggttaacg 18240
catgcttaca aaaccttata tatagaaatc gttgctgcat gcttatatgt ctgtgtttgt 18300
cattgtctag gacttaagtc tgaagagata caccaatag gtggttaggt tttgtatggt 18360
aattttgta ttgccatcca aaacaggcct ctgaatttgt gtatttctat tattaacaac 18420
ctgatttttg cagctcttaa gttacgtatt aacaaagtaa aaacctgtaa aatccgaggc 18480
ttctgttcac gaaactcatc cggtttattc ctttgttctt gttctctcct atatcatgct 18540
tcatccatcc aacatcgcgc acctcgctaa ccaataataa actgaacaaa aaaaaaaaa 18600
tatgaaatac taggcccacat ccgacatcga gtggaaattt ggcttgcca agcttatagc 18660
caatatgcat cgcaccacaa cagcgccatc gccgcttaat ttagttaacca cctggtttat 18720
gtggatcgtc gagaaggtca aggtaaaatc tcaggtagc aggtgcgct tccagccgct 18780

```



-continued

---

gtgctgtgt ctctctctct ctatccgtat cctgtatcct gtatcttata cctgtttcca 18840  
 tatctgttga ctatataaag tgcaactacc agaaccgata ctgaacgggt gtagtttget 18900  
 gaccttttcc ccaaccatt taaagcaatt tggcaacaac cgcaatgagt ttgaacacag 18960  
 tgaatgcttt aagtgtgttg cccacataag aaaatcacct tgcaccttg cactttctct 19020  
 gtaacttcaa aataggagat cgaatatag gtatgtaaat gtttcgatcc cctacactgt 19080  
 atggcacttt atgtccagca cttggcacc c gattgctttc gatgtaatga acatttgctg 19140  
 actgcttata tgttgtgtct cttgtcttgt atgtgatcta tgtcccggt ctaatgcgcc 19200  
 ttgatctaac ccacaaaacc tgcaaaaaa tcttgcaaac cgcaattcaa aaaacacgag 19260  
 cctcaggcac gcatgaagaa aaagaagcgt ccaagtctgg ttcagatgat gggaaacgt 19320  
 caggccagtc cgcgtaccaa agccggcgcc aagtggctgt cgaagatcaa gaaaggtgag 19380  
 acatgtatgt atcgtctgtg ggctactccg accaggatcc gtccatatcc tggaaaacac 19440  
 aaccatcca tccgaggggt tttgtagcta acagcgtgtc agcccaagtg taactcctaa 19500  
 ctttccttea actcaactct tttctctgga acaattggct cgctctagct cgaattatt 19560  
 tctcaacct ttcgccttcc cagtgcacaa aggtagaaac gccatggatc tctataaatc 19620  
 cgacattata ttgaatttga ggtagaagtc gtgatctttg gcgtttgtac ctcagtgcac 19680  
 cttgctgtat agtggaaatc aaaagctaat gatattacct cgaattccca gactcagtgg 19740  
 cctgtcgcga ggtccatcta tcgctcttgg gatcacaggc gagcttctcg caggccaatc 19800  
 agaatcgcac cgagaacgtg gccgactggg aggcgattgc caaaaagtac cgggactgg 19860  
 ttggcgacga ggagggtgga tcgctcaagg actcggatgc ggagagtgga tcgcaggagg 19920  
 gtagcggagg acaacagcca ccggcacagg tgggcagacg agccatcaag gccaccctgg 19980  
 cagacactac aaaatagaca cacagaaatg acacagaaaa aacagaaaaa cagcttcgga 20040  
 tgcttaatta actacgtttt gattgcaggt ctaagcttca tctatctctt caaactatcc 20100  
 ttctgacta tctctatctc ttctcgacta tccaagcgtc tgtccttctg taattctaag 20160  
 atctaactct aagaaactct atccgtaagc tgcaccttgg gtatggtttt ctcagactct 20220  
 ggaaccact tcttttgggt caactggagt atgggaaat cagactaaaa tcttaagtt 20280  
 aagccttcc tttctaaact aattttagct agaatttga aattgttttg agtaaccttt 20340  
 aaagcgaag ctgattgttt attttgatg gattttccgt tggagttttc tacgattagc 20400  
 gaaacaaca aaaaaagttt tccatgttcc agatttttaa agtaagttaa ttcgtccttt 20460  
 ttggactcaa tttgccttac atttttgaa accaactcct agcattttgt attaagctaa 20520  
 tgattgcgac catatcgta ataagtattg tcttagagat gttaaagtaa ttgaacttta 20580  
 gtttcaatcg gagctaaaag tcaagcgggt ttatataaat ctgcataat ctcattgttt 20640  
 tccgtaatt gtcaagtaac aacgttccct ctacttacta agctttgggt cattttttat 20700  
 aacaaatgag cgcataaaat tgttaactgt acttgattgt aaataaataa gtcttatttt 20760  
 aaaaattgt actattgctt cagcttgtaa tcattgcata ctttttggcg gcaactggcat 20820  
 ataccgccat ctatcggagg aaacaaaatt ttaaaattat gtttagcatt attttttcta 20880  
 attaaactat ttttgggttc atgcttataa tacaattata attttataat tataagtctg 20940  
 tatttttgaa taaatggatt gtttttgtgt ttgttattta tategtacgt tactcggctg 21000  
 ctgocagatt atcaaaaata gctctogctt atttccatt cacttgagcg acatctgtga 21060  
 atgaaatata gaacatgcgg ataaggtatt ttttggtttt cattaaatcc cgctaggtgg 21120

-continued

---

cgaatgcaaa	tgtaaaatta	atgtaaatg	ataaatcatt	gaaactaatg	attaaaaaaaa	21180
attgatttag	aatttaataa	tatatattgt	atthtgaata	atatttccta	aacctttcat	21240
ttaaataaaa	atgattacga	ttttatcata	aatgttggtt	tttattctaa	cttagtaact	21300
gcaagctggt	ttgattatgc	caagataatt	tcaaaatagg	ctagaattct	ctcctttaa	21360
ccatgtaatc	atggccataa	agctaagaac	gggcaataaa	atcgcttaa	ttgcctgct	21420
gaattgacag	attaccaaga	ggcactcagg	cgtcattagc	cgggccagca	gaaaagcgac	21480
agaaaccgca	tcggaatg	accaaggtgt	tgaacttcgg	aattgcattt	taatttggt	21540
tcaagctgca	gtttgctgtt	gttttcgct	cgattgcagg	tgtcacagtc	ggtttaaattg	21600
tgttgaaaac	ctcaagtggt	caatgtttgc	tgcttgctgc	actcgactc	gtattattac	21660
acataattgc	cccttgccgt	tgacattggt	gctgtgtggc	agttgcactt	gcatttgctg	21720
ttgctgctgt	gcttgatatt	tgccaccgat	aaaatgcata	catacatgca	aaaatatatg	21780
aaaaagcaaa	gcaaacgagt	ttctgtagcc	gcagccaagg	tttatggcca	caagcgtgtc	21840
aatttaagct	gcaattagc	agttaataaa	tttaaccgat	cttaccagtc	agataccagg	21900
tccagatgcc	agctgattaa	tgccacttcc	ccagcgattc	ggtagctgca	acgtacaaaa	21960
ctccaaatgg	attccaatcg	gattcgatgc	tggcgatgct	gtggctgtcc	gtcatccatc	22020
aaaggtttct	tctacggacc	aggaagcagt	ttcgattcga	ttcgatccgg	gcttccatgg	22080
cttcagcctc	cgcgactcgg	catcgtgcaa	catgtgtgtg	gtgtgttggc	acagcaggtg	22140
acatttccag	gccagatcag	gaaaatgtaa	ataaatgatc	ggacattgga	cgacacccat	22200
gcccataacc	ataccatata	gatcaacctg	gctgaacacg	acatggagca	agttgtacct	22260
ggttatacga	ctatatgttg	ctgttcatgt	tgctgttctg	ttgatataca	aaacactttt	22320
tcatatcgaa	atthtgtgata	ggccgtgatt	aatggcgagc	gacacaaaa	cttaatttga	22380
cgccaggccc	gtagctggcg	ccttggggaa	atggcagaga	tccgaacgca	aactctttgg	22440
gtgcacagag	agaaaagatg	ctaattttcc	attaaaagta	tttagtatca	gcttgaatga	22500
taggtaggtt	actgttaaag	cgthtctgtt	gagctaatag	gcattaataa	atgccattga	22560
acaactaaca	tttaagacta	ttttataagt	aatgagatca	taaatagtaa	aaatgtagtt	22620
acctcttttt	ttcatcctgt	agccttgaat	ttgctgctgg	tttctggct	gggagaaaa	22680
ataatctcgg	gcaagattaa	ttattgtaat	cacatcaaca	gcagagccat	gcaaacggat	22740
tctcgtatcc	gtattcgttt	tcgthtccgg	aatgggagtc	acagaaaaac	caacacgaaa	22800
atgatcaatg	atcatcctgt	ggttctctgt	tgatttttat	agcgaacgcc	cgatcgcctg	22860
cctgggttac	acatttcatt	ggctaataca	gatgctaatt	tgaagaagat	taattcgtgt	22920
gcgagtttct	gactgcctgc	caggcaagcc	cgaagattcg	aagattataa	tctgctaagc	22980
aagaggaaac	tgaaggctta	ttattaatac	aggccaacac	agccccaga	aatgtgtcct	23040
gtatttaatt	aaatacgcgc	acactgggaa	aagcaattcc	aatgaattct	taatctattt	23100
tctaatttta	taggacatta	aaaccatata	ttaaataaaa	aactcttgta	tcgaaatcat	23160
taaaatgta	tgcttacttg	caaagactta	tcaccatttt	tttcgctgt	atctgccatt	23220
tagccacatc	ccagaaatgt	ggagagtttc	gggtgagttg	tgcttggca	gtgcagtgac	23280
acgcagatta	attgaaatth	tatgagtagc	gcagacgtaa	acaatcagcg	agaccacctt	23340
ttgcagccc	cttaggtcat	aggagctcgc	caagatcccc	ctgctcggat	ggcgtatcca	23400
tgtccagatt	ccaagctcca	gcttgactac	actaactggc	caagtcggca	acggacagct	23460
gtggctcacc	ccgtggccaa	aagaaacttg	caacattatg	aaaatggac	cacagccatg	23520

-continued

---

```

cacagtgggtt gacagcagac ccttgggatg tgtggaaatt atttggaagc aacagcaaat 23580
aattccagat aatgcaatta attcgatact tatatattat attctatatg tatttttagta 23640
ttttaaagaa cttctgttga taccactgtg ccctgtgatc ctgctgacgc gatcgccacg 23700
ctaattgata gactgtgaaa ttatttaaca acggctggaa agtgagctcg gcgtggctgc 23760
ggctcgaaag gagcttccaa gcgtggccag atgggtcaga aggctttcga cccggccatc 23820
aagaccaggg tcggcacatc tttttggtgg ctctggtccc tggccgctgg ccaatcatcc 23880
atccagtgga ggatcgcgga cttacggcta agtgaaaagt gttaaaaagc acgactcacg 23940
gcgggcagtt gtgtcggatt tgaagacaaa tgagcagcgt cttttgacat ttgcgaaatt 24000
taaaatgtca gccgaaaact ggtgggtcgt ccacccttga cgaaggtttc ggatgggagg 24060
tcccggttcc atageggatc gccacgcttt gccggataag tcgcgagaaa tttaaattaa 24120
aactcaggtg aaaggttatt aattcgcaag tggaactggg gcgtagctcg gctcactggt 24180
aatactcgaa atctccactc atttgggta atgctgatgg cactttgaca gggatgatga 24240
tgatggggat atgacgaatg ccagcggcga tgatgccaaa taaaatggaa gtgacagagt 24300
tcagtgcggtt ggttttaatt aataagcata tttccagaga gctttctttt cagcaaaag 24358

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1704
<212> TYPE: PRT
<213> ORGANISM: Drosophila melanogaster
<220> FEATURE:
<223> OTHER INFORMATION: amino acid sequence derived from nompC genomic
sequence

```

```

<400> SEQUENCE: 2

```

```

Arg Thr Pro Met His Leu Ala Ala Glu Asn Gly His Ala His Val Ile
 1           5           10          15
Glu Ile Leu Ala Asp Lys Phe Lys Ala Ser Ile Phe Glu Arg Thr Lys
 20          25          30
Asp Gly Ser Thr Leu Met His Ile Ala Ser Leu Asn Gly His Ala Glu
 35          40          45
Cys Ala Thr Met Leu Phe Lys Lys Gly Val Tyr Leu His Met Pro Asn
 50          55          60
Lys Asp Gly Ala Arg Ser Ile His Thr Ala Ala Tyr Gly His Thr
 65          70          75          80
Gly Ile Ile Asn Thr Leu Leu Gln Lys Gly Glu Lys Val Asp Val Thr
 85          90          95
Thr Asn Asn Tyr Thr Ala Leu His Ile Ala Val Glu Ser Ala Lys Pro
100         105         110
Ala Val Val Glu Thr Leu Leu Gly Phe Gly Ala Asp Val His Val Arg
115         120         125
Gly Gly Lys Leu Arg Glu Thr Pro Leu His Ile Ala Ala Arg Val Lys
130         135         140
Asp Gly Asp Arg Cys Ala Leu Met Leu Leu Lys Ser Gly Ala Ser Pro
145         150         155         160
Asn Leu Thr Thr Asp Asp Cys Leu Thr Pro Val His Val Ala Ala Arg
165         170         175
His Gly Asn Leu Ala Thr Leu Met Gln Leu Leu Glu Asp Glu Gly Asp
180         185         190
Pro Leu Tyr Lys Ser Asn Thr Gly Glu Thr Pro Leu His Met Ala Cys
195         200         205

```

-continued

Arg	Ala	Cys	His	Pro	Asp	Ile	Val	Arg	His	Leu	Ile	Glu	Thr	Val	Lys
210						215					220				
Glu	Lys	His	Gly	Pro	Asp	Lys	Ala	Thr	Thr	Tyr	Ile	Asn	Ser	Val	Asn
225					230					235					240
Glu	Asp	Gly	Ala	Thr	Ala	Leu	His	Tyr	Thr	Cys	Gln	Ile	Thr	Lys	Glu
				245					250					255	
Glu	Val	Lys	Ile	Pro	Glu	Ser	Asp	Lys	Gln	Ile	Val	Arg	Met	Leu	Leu
			260					265					270		
Glu	Asn	Gly	Ala	Asp	Val	Thr	Leu	Gln	Thr	Lys	Thr	Ala	Leu	Glu	Thr
		275					280					285			
Ala	Phe	His	Tyr	Cys	Ala	Val	Ala	Gly	Asn	Asn	Asp	Val	Leu	Met	Glu
	290					295					300				
Met	Ile	Ser	His	Met	Asn	Pro	Thr	Asp	Ile	Gln	Lys	Ala	Met	Asn	Arg
305					310					315					320
Gln	Ser	Ser	Val	Gly	Trp	Thr	Pro	Leu	Leu	Ile	Ala	Cys	His	Arg	Gly
				325					330					335	
His	Met	Glu	Leu	Val	Asn	Asn	Leu	Leu	Ala	Asn	His	Ala	Arg	Val	Asp
			340					345					350		
Val	Phe	Asp	Thr	Glu	Gly	Arg	Ser	Ala	Leu	His	Leu	Ala	Ala	Glu	Arg
		355					360					365			
Gly	Tyr	Leu	His	Val	Cys	Asp	Ala	Leu	Leu	Thr	Asn	Lys	Ala	Phe	Ile
	370					375					380				
Asn	Ser	Lys	Ser	Arg	Val	Gly	Arg	Thr	Ala	Leu	His	Leu	Ala	Ala	Met
385					390					395					400
Asn	Gly	Phe	Thr	His	Leu	Val	Lys	Phe	Leu	Ile	Lys	Asp	His	Asn	Ala
				405					410					415	
Val	Ile	Asp	Ile	Leu	Thr	Leu	Arg	Lys	Gln	Thr	Pro	Leu	His	Leu	Ala
			420					425					430		
Ala	Ala	Ser	Gly	Gln	Met	Glu	Val	Cys	Gln	Leu	Leu	Leu	Glu	Leu	Gly
		435					440					445			
Ala	Asn	Ile	Asp	Ala	Thr	Asp	Asp	Leu	Gly	Gln	Lys	Pro	Ile	His	Val
	450					455					460				
Ala	Ala	Gln	Asn	Asn	Tyr	Ser	Glu	Val	Ala	Lys	Leu	Phe	Leu	Gln	Gln
465					470					475					480
His	Pro	Ser	Leu	Val	Asn	Ala	Thr	Ser	Lys	Asp	Gly	Asn	Thr	Cys	Ala
				485					490					495	
His	Ile	Ala	Ala	Met	Gln	Gly	Ser	Val	Lys	Val	Ile	Glu	Glu	Leu	Met
		500						505					510		
Lys	Phe	Asp	Arg	Ser	Gly	Val	Ile	Ser	Ala	Arg	Asn	Lys	Leu	Thr	Asp
		515					520					525			
Ala	Thr	Pro	Leu	Gln	Leu	Ala	Ala	Glu	Gly	Gly	His	Ala	Asp	Val	Val
	530					535					540				
Lys	Ala	Leu	Val	Arg	Ala	Gly	Ala	Ser	Cys	Thr	Glu	Glu	Asn	Lys	Ala
545					550					555					560
Gly	Phe	Thr	Ala	Val	His	Leu	Ala	Ala	Gln	Asn	Gly	His	Gly	Gln	Val
				565					570					575	
Leu	Asp	Val	Leu	Lys	Ser	Thr	Asn	Ser	Leu	Arg	Ile	Asn	Ser	Lys	Lys
			580					585					590		
Leu	Gly	Leu	Thr	Pro	Leu	His	Val	Ala	Ala	Tyr	Tyr	Gly	Gln	Ala	Asp
		595					600					605			
Thr	Val	Arg	Glu	Leu	Leu	Thr	Ser	Val	Pro	Ala	Thr	Val	Lys	Ser	Glu
	610					615					620				
Thr	Pro	Thr	Gly	Gln	Ser	Leu	Phe	Gly	Asp	Leu	Gly	Thr	Glu	Ser	Gly



-continued

---

Leu Leu Lys Asp Leu Ala Arg Phe Leu Ala Val Leu Ala Ile Phe Val  
 1060 1065 1070  
 Phe Gly Phe Ser Met His Ile Val Ala Leu Asn Gln Ser Phe Ala Asn  
 1075 1080 1085  
 Phe Ser Pro Glu Asp Leu Arg Ser Phe Glu Lys Lys Asn Arg Asn Arg  
 1090 1095 1100  
 Gly Tyr Phe Ser Asp Met Glu Gln Met Thr Cys Pro His Pro Asp Leu  
 1105 1110 1115 1120  
 Arg Arg Trp Arg Ile Met Ser Ile Val Ala Ser Ala Asn Ser Asp Glu  
 1125 1130 1135  
 Ser Thr Arg Thr Thr Phe Pro Gly Gly Thr Ser Thr Ser Pro His Ser  
 1140 1145 1150  
 Leu Leu Glu Ile Pro Ser Pro Cys Met His Val Asp Val Phe Ile Gln  
 1155 1160 1165  
 Ser Ile Gln Thr Lys Ile Lys Gln Ser Ile Ser Asn Ile Asp Ile Thr  
 1170 1175 1180  
 Asn Ala Arg Leu Pro Gly Ala Phe Ser Leu Arg Arg Leu Pro Thr Thr  
 1185 1190 1195 1200  
 Lys Phe Cys Thr Ile Glu Thr Ile Glu Thr Asp Arg Ile Glu Ser Ile  
 1205 1210 1215  
 Thr Lys Asn Asp Asn Ala Thr Asp Thr Asp Tyr Arg Cys Ser Tyr Met  
 1220 1225 1230  
 Leu Gly Pro Met Thr Pro Phe Leu Ala Phe Glu Arg Leu Phe Phe Ala  
 1235 1240 1245  
 Val Phe Gly Gln Thr Thr Thr Leu Asp Ile Asn Pro Met Arg His Leu  
 1250 1255 1260  
 Arg Pro Glu Trp Thr Glu Val Leu Phe Lys Phe Val Phe Gly Ile Tyr  
 1265 1270 1275 1280  
 Leu Leu Val Ser Val Val Val Leu Ile Asn Leu Leu Ile Ala Met Met  
 1285 1290 1295  
 Ser Asp Thr Tyr Gln Arg Ile Gln Met Asn Arg Asn Trp Gly Leu Val  
 1300 1305 1310  
 Asp Arg Thr Asn Gln Arg Asn Lys Lys Lys Lys Asn His Ile Ile  
 1315 1320 1325  
 Glu Ser Thr Asn Pro Thr Trp Ala Ser Val Ile Phe Leu Phe Phe Lys  
 1330 1335 1340  
 Ile Ile Ser Thr Pro Ala Asn Ile Cys Val Leu Ser Gly Gly Val Tyr  
 1345 1350 1355 1360  
 Leu Tyr Leu Tyr Leu Tyr Leu Glu Met Tyr Leu Trp Val Ser Asp Thr  
 1365 1370 1375  
 Val Arg Met His Pro Ile Asn Ser Phe Glu Leu Leu Phe Phe Ala Val  
 1380 1385 1390  
 Phe Gly Gln Thr Thr Thr Glu Gln Thr Gln Val Asp Lys Ile Lys Asn  
 1395 1400 1405  
 Val Ala Thr Pro Thr Gln Pro Tyr Trp Val Glu Tyr Leu Phe Lys Ile  
 1410 1415 1420  
 Val Phe Gly Ile Tyr Met Leu Val Ser Val Val Val Leu Ile Asn Leu  
 1425 1430 1435 1440  
 Leu Ile Ala Met Met Ser Asp Thr Tyr Gln Arg Ile Gln Ala Gln Ser  
 1445 1450 1455  
 Asp Ile Glu Trp Lys Phe Gly Leu Ser Lys Leu Ile Arg Asn Met His  
 1460 1465 1470

-continued

---

Arg Thr Thr Thr Ala Pro Ser Pro Leu Asn Leu Val Thr Thr Trp Phe  
           1475                                  1480                                  1485  
  
 Met Trp Ile Val Glu Lys Val Lys Val Lys Ser Gln Val Thr Lys Val  
           1490                                  1495                                  1500  
  
 Ala Phe Gln Pro Leu Ser Leu Cys Leu Ser Leu Ser Ile Arg Ile Leu  
           1505                                  1510                                  1515                                  1520  
  
 Tyr Pro Val Ser Tyr Thr Cys Phe His Ile Cys Met Lys Lys Lys Lys  
                                   1525                                  1530                                  1535  
  
 Arg Pro Ser Leu Val Gln Met Met Gly Ile Arg Gln Ala Ser Pro Arg  
                                   1540                                  1545                                  1550  
  
 Thr Lys Ala Gly Ala Lys Trp Leu Ser Lys Ile Lys Lys Ser Val Ala  
           1555                                  1560                                  1565  
  
 Leu Ser Gln Val His Leu Ser Pro Leu Gly Ser Gln Ala Ser Phe Ser  
           1570                                  1575                                  1580  
  
 Gln Ala Asn Gln Asn Arg Ile Glu Asn Val Ala Asp Trp Glu Ala Ile  
           1585                                  1590                                  1595                                  1600  
  
 Ala Lys Lys Tyr Arg Ala Leu Val Gly Asp Glu Glu Gly Gly Ser Leu  
                                   1605                                  1610                                  1615  
  
 Lys Asp Ser Asp Ala Glu Ser Gly Ser Gln Glu Gly Ser Gly Gly Gln  
           1620                                  1625                                  1630  
  
 Gln Pro Pro Ala Gln Val Gly Arg Arg Ala Ile Lys Ala Thr Leu Ala  
           1635                                  1640                                  1645  
  
 Asp Thr Thr Lys Ser Lys Leu His Leu Ser Leu Gln Thr Ile Leu Pro  
           1650                                  1655                                  1660  
  
 Asp Tyr Leu Tyr Leu Phe Ser Thr Ile Gln Ala Ser Val Leu Leu Cys  
           1665                                  1670                                  1675                                  1680  
  
 Thr Leu Gly Met Val Phe Ser Asp Ser Gly Thr His Phe Phe Trp Phe  
                                   1685                                  1690                                  1695  
  
 Asn Trp Ser Met Gly Lys Ser Asp  
           1700

<210> SEQ ID NO 3  
 <211> LENGTH: 6156  
 <212> TYPE: DNA  
 <213> ORGANISM: Drosophila melanogaster  
 <220> FEATURE:  
 <223> OTHER INFORMATION: nompC cDNA sequence

<400> SEQUENCE: 3

```

tttctcgtcg ctccgaaaaa aggcaaaata gtaggcaacc tgaatccag agttgtagtt      60
ggggactctt ttggccaaaa tacaaggagg agaaaaatag aaaataataa agggggcacc      120
gccgttaacg cacacgcaac cgaagccata aaggggctaa acatataaat ttgtgtagta      180
aaagtgaaga aagcgaaaga atcaaagtgg aataatagcg agtgtttttc ggtttgctag      240
tgtgtttctg agtcggagtt tgtgtgtgtg tgtttgtgtg attcctagtg tgtctgttgc      300
tgttgccaat gaaaatgcaa attgttggtg acaaatattg gtaaatgcg gaggccgtag      360
gaatttgtgc aatgcgagtg cgaagtgaag gagcccgaag ctatgcagct aaaaaccgc      420
cctcctacc cgcacgcaat caataataat acaataaccc aaacgtatta cacggataat      480
ggcagcataa accagttaac atccgacagt gtttccgect aaccatcgag cacctagctc      540
atccccctg ccaccaaccc ttcgaaaaat ccccatgatc agcgccggat tgtgggagcag      600
taactagcga gccataaccg gatgtgcgag ccgcgcgag ggcgtggcgg tgggcgtggc      660
ggcggagtgg gtcgcaaaac cccctcctcg ctgaccggcc caccggatga gtcggctacg      720
  
```

-continued

---

cccagcgaac	gggctacgcc	cgccagcaaa	gcagactccg	atcccaagga	cgatagctcg	780
agcaatggcg	acaagaagga	tatggatctt	tttccagccc	caaagccgcc	gagtgccggc	840
gcctccattc	gggacacggc	gaacaagggtg	ctcggattgg	ccatgaaaag	cgagtggacg	900
cccctcgagg	cgagactcaa	gaagctggaa	aagtatgtgg	ccaatgtggg	cgaggatggc	960
aatcacatac	cgctggccgg	cgttcacgac	atgaataccg	gcatgacgcc	gctgatgtac	1020
gcaacgaagg	acaataagac	ggccataatg	gatcgcatga	ttgagctggg	cgccgatgtg	1080
ggagcccgca	ataatgataa	ttataatgtg	ctacatattg	ccgcaatgta	ttcgcgtgag	1140
gatgtcgtca	aattgtgtct	aacaaaacgc	ggcgtggatc	ccttctccac	cggtggctcg	1200
cgttcgcaaa	ctgcggtgca	tttggtgtcc	agtcgacaaa	ccggaactgc	aactaatatc	1260
ctgcgcgctc	tgctcggcgc	agctggcaag	gatattcgct	tgaaagcggg	cggccgtggc	1320
aaaataccat	tgctcctggc	cgtggagtcg	ggcaaccagt	ccatgtgcag	ggagctcctg	1380
gctgcacaaa	cagcagagca	gctcaaggca	acgacggcca	atggagacac	ggccttgcat	1440
ttggccgcca	gacggcggga	cgtggacatg	gtccgcatcc	tggttgatta	cggaacgaat	1500
gtggacacgc	agaatgggga	gggccagacg	ccacttcata	tcgcggcgcg	cgaaggcgat	1560
gaggctctac	tcaagtactt	ctatggcgtg	cgcgcctcag	cgtccattgc	ggacaatcaa	1620
gatgcgactc	cgatgcactt	ggccgccgag	aatgggcacg	cgcacgtcat	cgagatactg	1680
gccgacaagt	tcaaggcgag	catcttcgag	cgcaccaagg	atggcagcac	gctgatgcac	1740
attgctgcac	tcaacggtea	tgctgagtcg	gccacgatgc	tcttcaagaa	ggcgtcttac	1800
ctccatagc	ccaacaagga	tggagcccgg	agtattcaca	ccgccgccgc	ctatggtcac	1860
acgggaatca	tcaacaccct	gctacagaag	ggcgagaaag	tggatgtgac	caccaatgat	1920
aactatacac	cactgcacat	agccgtggaa	tcggctaagc	ccgccgttgt	ggaaaccctg	1980
ctgggatttg	gagcagatgt	ccatgtccgt	ggcggaaaac	tacgtgagac	cccgtgcac	2040
attgccggac	gagtgaagga	tggagatagg	tgtgccctca	tgttgctgaa	gtcgggagcc	2100
agtccaaatt	tgaccacgga	tgactgtctg	acccccgtgc	atgtggcggc	tcgtcatggc	2160
aatctggcca	cgttgatgca	actcctcgag	gacgaaggag	atccgctgta	caaatcgaat	2220
actggagaga	caccgctgca	catggcctgt	cgtgcttgcc	acccggatat	tgtgcgtcat	2280
ctcatcgaga	cggatgaagga	gaaacacggt	cggataaagg	ccaccaccta	tataaactcg	2340
gtaaacgagg	acggcgccac	ggcgttgcat	tacacctgcc	aaatcaccaa	ggaggagggt	2400
aagattcccc	aatccgacaa	gcagatcggt	cggatgctcc	tcgaaaatgg	tcgggatgtc	2460
acgttgcaaa	cgaaaactgc	cttgagagacc	gctttccact	actgcgcggt	ggccggcaac	2520
aatgatgtgc	tgatggagat	gatctcacat	atgaatccca	cagacatcca	aaaggccatg	2580
aaccggcaat	catcgggtggg	ctggactcca	ctgctgattg	cttgccatcg	agggcacatg	2640
gagctggtea	ataatctact	ggcgaatcac	gctcgagtgg	atgtcttoga	tacggaagga	2700
cgatctgcct	tgcaattggc	tgctgagcga	ggatacctgc	atgtgtgtga	tgccctgctg	2760
accaataaag	cttttattaa	ctccaagtcc	cgcgtgggac	gcactgcact	acatctggca	2820
gccatgaatg	gatttacgca	tctggtgaaa	ttcctgatca	aggatcacia	tgcagttatc	2880
gatattctaa	cgttgagaaa	gcaaacgcgc	ctccatttgg	cggcagccag	cgggcagatg	2940
gaagtctgtc	agctgctcct	cgagctgggc	gccaatatcg	atgcgacgga	cgatctgggc	3000
cagaagccaa	tccacgtcgc	cgcccagaac	aactactctg	aagtggccaa	actcttctcg	3060
cagcagcatc	catccctggt	gaatgccacc	agcaaggatg	gaaacacatg	tgcccacatt	3120



-continued

---

gccgccatgc	agggatccgt	caaggtgatc	gaggagctga	tgaagttcga	tcgatcgggt	3180
gtgatttcgg	cgcggaataa	acttacggat	gccacgcccc	ttcagctggc	cgccgagggc	3240
ggacatgcgg	atgtggtgaa	ggctcctgtg	agagctggtg	cctcctgcac	cgaagagaac	3300
aaggcgggat	tcaccgccgt	tcattctggc	gcacagaatg	gacatggtca	ggtcctggat	3360
gtgctgaaaa	gcacaaactc	actaaggatc	aatagcaaaa	agttgggtct	gacgccgctt	3420
catgtggctg	cctattacgg	acaggcggat	accgtgcggg	aattgctgac	cagtgttccc	3480
gccaccgtca	agtcggaaac	tccaacggga	caaagtttat	ttggggatct	gggcacggag	3540
tccggaatga	caccactaca	cttggcggcc	ttttccggca	acgagaacgt	ggtgcgactg	3600
ctcctcaact	ctgcggtgtg	tcaagtggat	gcggcgacca	tcgagaacgg	ctataatcca	3660
ctccatttgg	cttgettcgg	tggteacatg	tcagtggteg	gtttgctcct	aagtcggteg	3720
gcggaactcc	tccaatcgca	ggatcgtaac	ggcaggacgg	gcctgcatat	cgccgccatg	3780
catggccaca	tccagatggt	ggagattctg	ctcggccagg	gcgcggagat	caacgcaacc	3840
gatcggaaeg	gttggacgcc	actgcattgt	gctgccaagg	ctggccactt	ggaggtgggtg	3900
aagtgtctgt	gcgagcgggg	tgctctcgca	aaatcggaga	ccaactacgg	ttgcgccggc	3960
atltggttcg	ccgcctccga	gggacacaac	gaggtcctgc	ggtatctgat	gaacaaggag	4020
cacgacacct	acggcctgat	ggaggacaag	cgattcgtgt	acaacctgat	ggtggtgtcc	4080
aagaaccaca	acaacaagcc	cattcaggag	tttgtcctgg	tatcaccagc	accctgggat	4140
acagccgcca	aactgtccaa	catctacata	gtactctcga	caaaggaaaa	agagcgcgcc	4200
aaggatctgg	tagcagctgg	caaacagtgc	gaggcaatgg	ccacggagct	cttggccctg	4260
gcagctgggt	cagattccgc	cggaaagatc	cttcaagcca	ccgataagcg	aaacgtggag	4320
tttctcgacg	ttctcattga	aaatgagcag	aaggaagtga	ttgccacac	ggtagttcag	4380
cgatacttgc	aagaactctg	gcattggctcc	ctgacgtggg	catcctggaa	aatccttctg	4440
ctgctcgtgg	ccttcatagt	ctgcccacca	gtgtggattg	gattcacatt	cccgatgggt	4500
cacaagtcca	acaaggtgcc	catcatcaag	ttcatgtcgt	acctaacctc	tcacatttac	4560
ctcatgatcc	acctgagcat	cgtgggcata	acgcccattt	accagtgct	ccgattgagt	4620
ttggtgccct	actggtacga	ggtgggtctt	ctcatctggc	tgagtggatt	gctccttttc	4680
gagctgacga	atccgtcaga	taaactggga	ctgggatcga	taaagtgct	cgtgctgctg	4740
ctcggcatgg	ccggagtggt	tgtccatgtc	tcagcatttc	tattcgtctc	caaggagtac	4800
tggccaactt	tgggtgattg	tcgaaatcag	tgettccgct	tggccttcc	gctggcctgt	4860
gtgcagatcc	tcgacttttt	gtccttccac	cacctattcg	gtccctgggc	catcatcatt	4920
ggggatctgc	tgaaggatct	ggctcggttt	ttggccgtcc	tggccatctt	tgtgtttggc	4980
ttttccatgc	acattgtggc	cctgaaatcag	agctttgcca	atltctcacc	ggaggatctg	5040
cgcagcttcg	agaagaagaa	ccgaaataga	ggctacttca	gtgacgtgcg	catgcatccg	5100
attaactcgt	tcgagttggt	gttcttcgcc	gtgttcggac	aaacgacgac	cgagcaaacg	5160
caagttgaca	aaatcaaaaa	tgtagccacg	cccactcaac	cgtattgggt	tgagtacctg	5220
ttcaaaattg	tctttggcat	ttacatgttg	gtgtcgggtg	ttgtgctcat	taacctgctg	5280
attgctatga	tgtcagacac	ctatcaacgc	attcaggtag	tattgctaaa	tgcgctttta	5340
tctaactcga	ctctatttat	taactcgtac	tttaaccata	agtatataaa	tttcatattg	5400
cattgtgtat	taatcattct	ctatttcagc	ataagaagta	aatttacata	tgaagatgat	5460

-continued

---

```

ttatatttct tagatatata atagcggtag ttaggaagtg agctgttttg ggaacatatt 5520
gagaaaaatag ttaattaatc tggagaactt ggcatgctct gtaaatccat caactgccca 5580
gacttgcac cttccagggtt tttcaggaaa ataatgtag caatctgagg gatacaattt 5640
tgtgaaagtg tatctcaaag atggaagcct gccgccttct agtgtagtac agtgcagagt 5700
agcttttagtg gattagccgc cttgaagtgt gccctgcttt tgtgaccagt gttgagcgag 5760
gccaaaccag aaagtgttgg ttaacgcatg cttacaaaac cttatatata gaaatcgttg 5820
ctgcatgctt atatgtctgt gtttgcatt gtctaggact taagtctgaa gagatacacc 5880
aatatggtgg ttaggttttg tatggtaatt ttgtgattgc catccaaaac aggctctga 5940
atattgtgat ttctattatt aacaacctga tttttgcagc tcttaagtta cgtattaaca 6000
aagtaaaaaa ctgtaaaatc cgaggcttct gttcacgaaa ctcatcccgt ttattccttt 6060
gtttctgttc tctctatat catgtctcat ccatccaaca tcgcgcacct cgctaaccaa 6120
taataaactg aacaaaaaaaa aaaaaaaaaa actcga 6156

```

```

<210> SEQ ID NO 4
<211> LENGTH: 1619
<212> TYPE: PRT
<213> ORGANISM: Drosophila melanogaster
<220> FEATURE:
<223> OTHER INFORMATION: amino acid sequence derived from nompC cDNA
sequence

```

```

<400> SEQUENCE: 4

```

```

Met Ser Gln Pro Arg Gly Gly Arg Gly Gly Gly Arg Gly Gly Gly Val
  1          5          10          15
Gly Arg Lys Thr Pro Ser Ser Leu Thr Gly Pro Pro Asp Glu Ser Ala
          20          25          30
Thr Pro Ser Glu Arg Ala Thr Pro Ala Ser Lys Ala Asp Ser Asp Pro
          35          40          45
Lys Asp Asp Ser Ser Ser Asn Gly Asp Lys Lys Asp Met Asp Leu Phe
          50          55          60
Pro Ala Pro Lys Pro Pro Ser Ala Gly Ala Ser Ile Arg Asp Thr Ala
          65          70          75          80
Asn Lys Val Leu Gly Leu Ala Met Lys Ser Glu Trp Thr Pro Ile Glu
          85          90          95
Ala Glu Leu Lys Lys Leu Glu Lys Tyr Val Ala Asn Val Gly Glu Asp
          100          105          110
Gly Asn His Ile Pro Leu Ala Gly Val His Asp Met Asn Thr Gly Met
          115          120          125
Thr Pro Leu Met Tyr Ala Thr Lys Asp Asn Lys Thr Ala Ile Met Asp
          130          135          140
Arg Met Ile Glu Leu Gly Ala Asp Val Gly Ala Arg Asn Asn Asp Asn
          145          150          155          160
Tyr Asn Val Leu His Ile Ala Ala Met Tyr Ser Arg Glu Asp Val Val
          165          170          175
Lys Leu Leu Leu Thr Lys Arg Gly Val Asp Pro Phe Ser Thr Gly Gly
          180          185          190
Ser Arg Ser Gln Thr Ala Val His Leu Val Ser Ser Arg Gln Thr Gly
          195          200          205
Thr Ala Thr Asn Ile Leu Arg Ala Leu Leu Ala Ala Ala Gly Lys Asp
          210          215          220
Ile Arg Leu Lys Ala Asp Gly Arg Gly Lys Ile Pro Leu Leu Leu Ala
          225          230          235          240

```

-continued

---

Val Glu Ser Gly Asn Gln Ser Met Cys Arg Glu Leu Leu Ala Ala Gln  
 245 250 255  
 Thr Ala Glu Gln Leu Lys Ala Thr Thr Ala Asn Gly Asp Thr Ala Leu  
 260 265 270  
 His Leu Ala Ala Arg Arg Arg Asp Val His Met Val Arg Ile Leu Val  
 275 280 285  
 Asp Tyr Gly Thr Asn Val Asp Thr Gln Asn Gly Glu Gly Gln Thr Pro  
 290 295 300  
 Leu His Ile Ala Ala Ala Glu Gly Asp Glu Ala Leu Leu Lys Tyr Phe  
 305 310 315 320  
 Tyr Gly Val Arg Ala Ser Ala Ser Ile Ala Asp Asn Gln Asp Arg Thr  
 325 330 335  
 Pro Met His Leu Ala Ala Glu Asn Gly His Ala His Val Ile Glu Ile  
 340 345 350  
 Leu Ala Asp Lys Phe Lys Ala Ser Ile Phe Glu Arg Thr Lys Asp Gly  
 355 360 365  
 Ser Thr Leu Met His Ile Ala Ser Leu Asn Gly His Ala Glu Cys Ala  
 370 375 380  
 Thr Met Leu Phe Lys Lys Gly Val Tyr Leu His Met Pro Asn Lys Asp  
 385 390 395 400  
 Gly Ala Arg Ser Ile His Thr Ala Ala Tyr Gly His Thr Gly Ile  
 405 410 415  
 Ile Asn Thr Leu Leu Gln Lys Gly Glu Lys Val Asp Val Thr Thr Asn  
 420 425 430  
 Asp Asn Tyr Thr Ala Leu His Ile Ala Val Glu Ser Ala Lys Pro Ala  
 435 440 445  
 Val Val Glu Thr Leu Leu Gly Phe Gly Ala Asp Val His Val Arg Gly  
 450 455 460  
 Gly Lys Leu Arg Glu Thr Pro Leu His Ile Ala Ala Arg Val Lys Asp  
 465 470 475 480  
 Gly Asp Arg Cys Ala Leu Met Leu Leu Lys Ser Gly Ala Ser Pro Asn  
 485 490 495  
 Leu Thr Thr Asp Asp Cys Leu Thr Pro Val His Val Ala Ala Arg His  
 500 505 510  
 Gly Asn Leu Ala Thr Leu Met Gln Leu Leu Glu Asp Glu Gly Asp Pro  
 515 520 525  
 Leu Tyr Lys Ser Asn Thr Gly Glu Thr Pro Leu His Met Ala Cys Arg  
 530 535 540  
 Ala Cys His Pro Asp Ile Val Arg His Leu Ile Glu Thr Val Lys Glu  
 545 550 555 560  
 Lys His Gly Pro Asp Lys Ala Thr Thr Tyr Ile Asn Ser Val Asn Glu  
 565 570 575  
 Asp Gly Ala Thr Ala Leu His Tyr Thr Cys Gln Ile Thr Lys Glu Glu  
 580 585 590  
 Val Lys Ile Pro Glu Ser Asp Lys Gln Ile Val Arg Met Leu Leu Glu  
 595 600 605  
 Asn Gly Ala Asp Val Thr Leu Gln Thr Lys Thr Ala Leu Glu Thr Ala  
 610 615 620  
 Phe His Tyr Cys Ala Val Ala Gly Asn Asn Asp Val Leu Met Glu Met  
 625 630 635 640  
 Ile Ser His Met Asn Pro Thr Asp Ile Gln Lys Ala Met Asn Arg Gln  
 645 650 655

-continued

---

Ser Ser Val Gly Trp Thr Pro Leu Leu Ile Ala Cys His Arg Gly His  
 660 665 670  
 Met Glu Leu Val Asn Asn Leu Leu Ala Asn His Ala Arg Val Asp Val  
 675 680 685  
 Phe Asp Thr Glu Gly Arg Ser Ala Leu His Leu Ala Ala Glu Arg Gly  
 690 695 700  
 Tyr Leu His Val Cys Asp Ala Leu Leu Thr Asn Lys Ala Phe Ile Asn  
 705 710 715 720  
 Ser Lys Ser Arg Val Gly Arg Thr Ala Leu His Leu Ala Ala Met Asn  
 725 730 735  
 Gly Phe Thr His Leu Val Lys Phe Leu Ile Lys Asp His Asn Ala Val  
 740 745 750  
 Ile Asp Ile Leu Thr Leu Arg Lys Gln Thr Pro Leu His Leu Ala Ala  
 755 760 765  
 Ala Ser Gly Gln Met Glu Val Cys Gln Leu Leu Leu Glu Leu Gly Ala  
 770 775 780  
 Asn Ile Asp Ala Thr Asp Asp Leu Gly Gln Lys Pro Ile His Val Ala  
 785 790 795 800  
 Ala Gln Asn Asn Tyr Ser Glu Val Ala Lys Leu Phe Leu Gln Gln His  
 805 810 815  
 Pro Ser Leu Val Asn Ala Thr Ser Lys Asp Gly Asn Thr Cys Ala His  
 820 825 830  
 Ile Ala Ala Met Gln Gly Ser Val Lys Val Ile Glu Glu Leu Met Lys  
 835 840 845  
 Phe Asp Arg Ser Gly Val Ile Ser Ala Arg Asn Lys Leu Thr Asp Ala  
 850 855 860  
 Thr Pro Leu Gln Leu Ala Ala Glu Gly Gly His Ala Asp Val Val Lys  
 865 870 875 880  
 Ala Leu Val Arg Ala Gly Ala Ser Cys Thr Glu Glu Asn Lys Ala Gly  
 885 890 895  
 Phe Thr Ala Val His Leu Ala Ala Gln Asn Gly His Gly Gln Val Leu  
 900 905 910  
 Asp Val Leu Lys Ser Thr Asn Ser Leu Arg Ile Asn Ser Lys Lys Leu  
 915 920 925  
 Gly Leu Thr Pro Leu His Val Ala Ala Tyr Tyr Gly Gln Ala Asp Thr  
 930 935 940  
 Val Arg Glu Leu Leu Thr Ser Val Pro Ala Thr Val Lys Ser Glu Thr  
 945 950 955 960  
 Pro Thr Gly Gln Ser Leu Phe Gly Asp Leu Gly Thr Glu Ser Gly Met  
 965 970 975  
 Thr Pro Leu His Leu Ala Ala Phe Ser Gly Asn Glu Asn Val Val Arg  
 980 985 990  
 Leu Leu Leu Asn Ser Ala Gly Val Gln Val Asp Ala Ala Thr Ile Glu  
 995 1000 1005  
 Asn Gly Tyr Asn Pro Leu His Leu Ala Cys Phe Gly Gly His Met Ser  
 1010 1015 1020  
 Val Val Gly Leu Leu Leu Ser Arg Ser Ala Glu Leu Leu Gln Ser Gln  
 1025 1030 1035 1040  
 Asp Arg Asn Gly Arg Thr Gly Leu His Ile Ala Ala Met His Gly His  
 1045 1050 1055  
 Ile Gln Met Val Glu Ile Leu Leu Gly Gln Gly Ala Glu Ile Asn Ala  
 1060 1065 1070  
 Thr Asp Arg Asn Gly Trp Thr Pro Leu His Cys Ala Ala Lys Ala Gly

-continued

1075			1080			1085									
His	Leu	Glu	Val	Val	Lys	Leu	Leu	Cys	Glu	Ala	Gly	Ala	Ser	Pro	Lys
1090					1095						1100				
Ser	Glu	Thr	Asn	Tyr	Gly	Cys	Ala	Ala	Ile	Trp	Phe	Ala	Ala	Ser	Glu
1105					1110						1115				1120
Gly	His	Asn	Glu	Val	Leu	Arg	Tyr	Leu	Met	Asn	Lys	Glu	His	Asp	Thr
					1125						1130				1135
Tyr	Gly	Leu	Met	Glu	Asp	Lys	Arg	Phe	Val	Tyr	Asn	Leu	Met	Val	Val
					1140										1150
Ser	Lys	Asn	His	Asn	Asn	Lys	Pro	Ile	Gln	Glu	Phe	Val	Leu	Val	Ser
Pro	Ala	Pro	Val	Asp	Thr	Ala	Ala	Lys	Leu	Ser	Asn	Ile	Tyr	Ile	Val
Leu	Ser	Thr	Lys	Glu	Lys	Glu	Arg	Ala	Lys	Asp	Leu	Val	Ala	Ala	Gly
1185					1190						1195				1200
Lys	Gln	Cys	Glu	Ala	Met	Ala	Thr	Glu	Leu	Leu	Ala	Leu	Ala	Ala	Gly
					1205						1210				1215
Ser	Asp	Ser	Ala	Gly	Lys	Ile	Leu	Gln	Ala	Thr	Asp	Lys	Arg	Asn	Val
					1220										
Glu	Phe	Leu	Asp	Val	Leu	Ile	Glu	Asn	Glu	Gln	Lys	Glu	Val	Ile	Ala
					1235										
His	Thr	Val	Val	Gln	Arg	Tyr	Leu	Gln	Glu	Leu	Trp	His	Gly	Ser	Leu
Thr	Trp	Ala	Ser	Trp	Lys	Ile	Leu	Leu	Leu	Val	Ala	Phe	Ile	Val	
1265					1270										1280
Cys	Pro	Pro	Val	Trp	Ile	Gly	Phe	Thr	Phe	Pro	Met	Gly	His	Lys	Phe
					1285										1295
Asn	Lys	Val	Pro	Ile	Ile	Lys	Phe	Met	Ser	Tyr	Leu	Thr	Ser	His	Ile
					1300										
Tyr	Leu	Met	Ile	His	Leu	Ser	Ile	Val	Gly	Ile	Thr	Pro	Ile	Tyr	Pro
					1315										
Val	Leu	Arg	Leu	Ser	Leu	Val	Pro	Tyr	Trp	Tyr	Glu	Val	Gly	Leu	Leu
					1330										
Ile	Trp	Leu	Ser	Gly	Leu	Leu	Leu	Phe	Glu	Leu	Thr	Asn	Pro	Ser	Asp
1345					1350										1360
Lys	Ser	Gly	Leu	Gly	Ser	Ile	Lys	Val	Leu	Val	Leu	Leu	Leu	Gly	Met
					1365										1375
Ala	Gly	Val	Gly	Val	His	Val	Ser	Ala	Phe	Leu	Phe	Val	Ser	Lys	Glu
					1380										
Tyr	Trp	Pro	Thr	Leu	Val	Tyr	Cys	Arg	Asn	Gln	Cys	Phe	Ala	Leu	Ala
					1395										
Phe	Leu	Leu	Ala	Cys	Val	Gln	Ile	Leu	Asp	Phe	Leu	Ser	Phe	His	His
					1410										
Leu	Phe	Gly	Pro	Trp	Ala	Ile	Ile	Ile	Gly	Asp	Leu	Leu	Lys	Asp	Leu
1425					1430										1440
Ala	Arg	Phe	Leu	Ala	Val	Leu	Ala	Ile	Phe	Val	Phe	Gly	Phe	Ser	Met
					1445										1455
His	Ile	Val	Ala	Leu	Asn	Gln	Ser	Phe	Ala	Asn	Phe	Ser	Pro	Glu	Asp
					1460										1470
Leu	Arg	Ser	Phe	Glu	Lys	Lys	Asn	Arg	Asn	Arg	Gly	Tyr	Phe	Ser	Asp
					1475										
Val	Arg	Met	His	Pro	Ile	Asn	Ser	Phe	Glu	Leu	Leu	Phe	Phe	Ala	Val
					1490										1500

-continued

Phe Gly Gln Thr Thr Thr Glu Gln Thr Gln Val Asp Lys Ile Lys Asn  
1505 1510 1515 1520

Val Ala Thr Pro Thr Gln Pro Tyr Trp Val Glu Tyr Leu Phe Lys Ile  
1525 1530 1535

Val Phe Gly Ile Tyr Met Leu Val Ser Val Val Val Leu Ile Asn Leu  
1540 1545 1550

Leu Ile Ala Met Met Ser Asp Thr Tyr Gln Arg Ile Gln Val Val Leu  
1555 1560 1565

Leu Asn Ala Leu Leu Ser Asn Ser Thr Leu Phe Ile Asn Ser Tyr Phe  
1570 1575 1580

Asn His Lys Tyr Ile Asn Phe Ile Leu His Cys Val Leu Ile Ile Leu  
1585 1590 1595 1600

Tyr Phe Ser Ile Arg Ser Lys Phe Thr Tyr Glu Asp Asp Leu Tyr Phe  
1605 1610 1615

Leu Asp Ile

<210> SEQ ID NO 5

<211> LENGTH: 9758

<212> TYPE: DNA

<213> ORGANISM: Caenorhabditis elegans

<220> FEATURE:

<223> OTHER INFORMATION: nompC genomic nucleotide sequence

<400> SEQUENCE: 5

```

ctttgccgct taaaattttg cagtgacata tccttatgga acactttcaa atgacacatg    60
tctcgtttta aagtctgacg gtaaactaaa aacatttcct tgtaagccta aacctaagcc    120
aaagcctaag cctaataagc ctagctaacg ctcgccactg acgccaagcc taagactaat    180
cctacgccaa tgccataaac tgacactgaa ataaaagtca aaagccaaaa gccaaaagcc    240
aaaaacctaag gccgaagcat aaggccaaag cctatgccta agcctgagcc tgagcttaaa    300
tcctaagcct aagcctaagg ccaaagaaca agcctaagtc taagtccaag cctaagtatc    360
aaaaacttac accgattccg ccaggctacc ctcagcacia ttatcaactt tgttaacata    420
tttatcggcg acggcggtgg gcttttctta ttcattctgc tgatcagaat agctcttccg    480
aacttccatt cttatccga ctgtgacctg attcgttggg aggtgtcaga catcatagca    540
atcagcaagt tgatcagcac aatcaagggt accatcatgt agattccgaa tagaagtttt    600
aagatgattt ttgcaaaatc tggaaactaga tggagcgggg gcattgaatc gggctcgacg    660
agtccgaaga gcgagaagaa gagcatttcg agggtttgag acggggaggg cagacgcatc    720
agctcggcgc tgtcctcctg gacaggctgg taggcaggct gaaaaaatct ctttcaaggg    780
tcgtttttct tgccataact acctggaaga tactcgtcac gtggagtgtg aagcccgcc    840
cgaacaacat caggatcaca aggaaacggg ccaaatcata cattagatcg ctgaaagcct    900
ctttctctaa ggggtcagct caagccaagt actcaccgaa taatgatcgc ccagggaccg    960
aacaatgat gcactgtcag gaaatccagg tactctacaa aagcaaatag cagggcaaag    1020
gcgaaaagtt gatttttcaa ataaagcatt gtcggggcga aatgtagctt ttcactgta    1080
tccaggtggg ttaggaatac tgccgggagc aggaaggcta ggacatggac ggctatcggc    1140
atcggggaaa ggactaggat taggaacttt acgattccta ggccagatcc tccaccgaca    1200
gtggagagtt cggagaccag atttccagag agccagagca acaggagcca ttccacaggg    1260
tttgaacca ccgaagttac ttcgtacctg gaaattgaga ttttgagggt ctatctgata    1320
tcccctaaat aaaatttaaa aaaataactt acatcttatg tgtaatatc aacaccacaa    1380

```

-continued

---

ttgtcagcag	tatcgtaaaa	tagacatgag	acacgatatg	gcacacaaat	ttaataatcg	1440
gagctcttcc	gatccgacta	tccagtggaa	gtgagaagta	gaacctatgcc	ggggggcata	1500
ttagcacgaa	gagggagaat	gcgacaaact	ttccgaatga	ccagtcgaca	cgggcagtcc	1560
atacttctgt	caggtagcgt	tggacagacg	cgtaggagac	tacttcttcc	tggaaacggg	1620
gctgttgagg	gttgactggt	taggttaagc	ttggagtgtt	acctgctcat	tttcaatgag	1680
aacatctagt	aggggcccgc	ctcgattgtc	cttagccttc	aggagaagag	cggcattgta	1740
ttcggtggeg	gtgatccctg	aaataatcta	ggactagtaa	attgtaagtc	attttctgaa	1800
aagattaaat	agctaagtgg	acctgtagcc	ttggccggta	actttggtcc	aataaccttg	1860
gtccagtaac	cttaatcctg	taaaccttgg	tcctgaaatc	ttggcctagt	aacctaaaac	1920
cttggtcctg	tggctcctgac	cctgttccctg	tatccttggg	tgggaaaccc	tagtccttgt	1980
cctggttttg	aaaccttggc	ccggtagcct	tggtcacagg	actggtcctg	tgccttggg	2040
cctggttctg	gtcttgggtc	cgaacacctg	gtccggcagt	tttggttctg	gtaccttggg	2100
cgtgtaacct	taaacccagt	aaccttggac	cggtaacctt	ggtacagtaa	ctttggtccg	2160
gaagccctgg	ctcggtaact	ctggtcctgg	tcattggtgt	ggtcctggcc	cggacaccct	2220
ggtccggtaa	ccctggtcta	gcaaccttgg	tcttgaccta	acaaccttgg	ttctgtaacc	2280
ttggtcttgt	aacttcggcc	ctgtatcctt	ggcccaaaga	ccttgggtccg	acagccttgg	2340
ttctgatacc	ttggtccagt	aactttggtc	gtggtcctgg	ttcaggtcca	gtaaccttga	2400
cccgataatc	ctggctctac	ctagtgaacct	tggcccggta	atcctgatcc	tggcccagta	2460
accttggtec	agtaecgtgg	ccctgcaact	atggcctagt	agctttggtc	cagtagccct	2520
gatcccgaat	ccttggttca	gtaaccttgg	tcttgggtcca	gtaactttgg	tctagtaacc	2580
atagtcacgt	aaccttggtc	ctgtaacctt	ggcccgctag	ccccgtagt	catgttcccg	2640
ctcctggtec	ggcagcattg	gtccggtaat	tttggacctc	ccctgggcct	tggcccaggg	2700
catgttctct	gtccaggggg	ccattttctt	cgtttttcat	tacctacctc	acaactccac	2760
agccatattc	tactgaaca	ctgccacatt	caacagatcc	ttcgccctct	ccttctcctt	2820
ctccgacata	tctctgtaca	acgcccagaa	cttgactgcc	gtctcaattg	gagcaggtga	2880
ttgaagaata	aactcttgta	gaggctcatt	gtcattggtt	ttaccacaaa	ccatcaagtc	2940
gaatatgaac	ttccgatcct	ccatcaattg	atgtgtgtca	tgcttctggt	tcaggaggaa	3000
tgaagacat	tctatatgat	tatgagctgc	agcaaagcac	aatggaactt	tgcctcctt	3060
ggtctccgcc	aatggatccg	ctgaactatc	gatgaacagc	ttgacgacac	tcaggtgcc	3120
ggcacgagtg	gcaaagttaa	gaccagtcca	gccattctga	tccatgacat	tgatgttaga	3180
tcctgagca	atgagaagtg	agaccatctc	gtagtggcca	ttctgagcgg	ctaggtggag	3240
cggggtcctg	cctctccaat	ccttggcgtg	ctgctgctga	gtagatctgg	acaggagcat	3300
tcctaccact	gcgatgtggc	cttgtggggc	agccagatgg	agggggatca	cgttctgaaa	3360
cggaaattha	aacgggggtca	ctgaaaaatt	caagttacca	ttgtagtact	ggtcgcgtca	3420
acttgactc	cctgattcag	aagcatccgc	acaagactgt	cgtgtccact	atgagcggct	3480
aaatggagag	gtgtgaagcc	gtattcagtt	gagaattcct	tattgacatg	gtgattgtag	3540
atgggcggct	cggaaacggac	tgttgcttgt	acgtgcttga	gcatttcatt	gacgaaatcc	3600
gaatttccgt	agaacgcagc	gatgtggaga	gcgttgagac	cggctcggaa	atgctaggtt	3660
cagggggaat	cgagtttttt	ttcagtacaa	aattcataaa	atttaaggct	agctgtgaaa	3720

-continued

---

aattgtgcta	ccaaagtata	ggccacggct	tcaaatttga	caggacttat	tccactttgc	3780
agatcagacc	tttatgcatg	aactgtactg	ccacgtattg	gaaaatgtta	tttttgacag	3840
ccttaccttt	ctcgaacacc	gtttccatag	gatcttatcg	aatgcctcca	aaatcgatat	3900
gaatccgttt	ttggcgccaa	ggtggagagc	agtcattccg	tgctgaaaat	caattctgcc	3960
taaaaatcgg	taaaagaacc	cctacccaat	tctcatcttc	cgcgtttgc	ccattctcca	4020
gcagaatcct	cacaatgttc	gcgtgacctc	ccgcagctgc	catatgaagt	gtagtggctt	4080
ccagtgtttt	ggtctttgcc	tggattacca	taggcttgtc	gatcatcata	agctcacgga	4140
ccacggctag	ggaaccttgg	aacaatatta	ttttagtgtc	aatcaaaagc	tgaagcttcc	4200
accctacctc	tcatcgacgc	aatatgtgcg	caggtgaatc	cattatgatc	aattgctggc	4260
aacacactcc	ggttggtatt	tctcattttc	aggaagagct	tcacaacgtc	ggggaagtca	4320
ttctcagctg	ccagatggag	aggggtttga	cccttgctcg	cacgtgcatt	ggggtttgct	4380
ccgagagcca	gaagggtttg	actcacagct	agctgaccga	attttgctgc	aaagtggagg	4440
gctgtctgga	aattatttgt	gtttctaata	aggagcttgc	cgacaaatth	gctcgaaccc	4500
cgtattagaa	actacgcaga	accctgtctg	ggcagtagat	tacctctagc	ttggatacta	4560
tcttacctga	ttatccagcg	taattgcctc	cagcgctgca	ccatgatcct	gcaccaggac	4620
attcaccacc	ttcacatgac	catgctgagc	tgctaagtgg	agcgggtcct	ctccggtttt	4680
cgatttactg	ttcacgaatg	ctttgtgctg	cagaagaagg	tgaaccaggg	agagatgccc	4740
attgaaagct	gccaggtgca	gagcagtacg	gcccatttca	tcgaatacat	caatacgggc	4800
gtggtgctga	aaaagtatga	ggtatccggt	ttgtgagaaa	tcagtggctc	cccagtagcc	4860
ttggcacagt	aaccttggtc	ctggtcctgg	tcctcgccca	gtaaccttgg	tcctgtaacc	4920
ctggtcctgt	agccctggcc	ctggtcctgg	tcctggctct	ggcccagtaa	ccttggtaact	4980
gtaaccatgg	tactgtaacc	ctggccctgg	tcctggctca	gtaacctcgg	ccttggtaacc	5040
ttggtcctgg	tcctggctct	ggtcctacac	acaaaaccag	taccttcaac	aaaatattcg	5100
ccactccaga	atgcctctg	gcacatgctt	ccaacagcgg	tgaccatccg	ttcttgcctc	5160
gcttgttctg	cacgatttgc	accgcaccgg	ctccgatctt	attgaccatc	gccaggagta	5220
cagcttgatt	tcggatcctt	gccgccat	gcacgcctg	ctcatttgc	ttgagtgatg	5280
gcatttctac	cattccaccg	tagtgatca	gaagatttac	tagcttggca	tcttctcctg	5340
gaaagtgtaa	ctggcgctgc	tcgatttcag	cggcgtagt	aagagctgtg	aagccgtcct	5400
gaaaaattta	acttgaagct	tcctgagatc	cagagaaaga	agctcacatt	ggttctatga	5460
ttgacatggt	ccttaagctg	ttcttgggtc	agaacttccg	aaaggctctt	caaaatcatt	5520
gatgctgctt	caaaattgca	tgacttggcg	gccacctgga	ggggtgtctc	tccgatcttt	5580
gagcttattt	tcgagtcgce	gttctcgtca	agcaggagcc	tggaaaaaag	gaggttcttg	5640
ggcttttaca	ggatccgaca	gaaaatagat	ttctcgaact	ttttccggtt	ttcgtactgt	5700
caatttacca	aatttcaagg	tacctgtttt	ttataagtgc	ttagaaatth	caaaaatttc	5760
aaaaattgtg	ataaactggg	gcgctgaatc	cagaattggc	acagaaatth	agagtttctc	5820
aattttcaaa	gaggcttgta	tgcaatgctt	agaaatceta	aattttgagc	acgcagttca	5880
cgggtccag	gaccaagtgc	acaataatct	caaaatthtt	gggtcccaca	gcagttgctc	5940
gctagctgaa	aaattctgca	cggcatgaga	agtggcacct	gtacgcaatt	tgtctaccgt	6000
atacctggac	gttttagtag	gtttttttca	aaattttttg	gaccaaaagct	tttttctca	6060
aaacgcgcct	aaacgtggct	aaactgcaat	tatcagttga	gcgcgtttac	actgatatac	6120



-continued

---

actttgcagg gccgtgtgct gattggctct aaagtcggcg tggctaagca ctgattagtc	6180
aagatcacct acttaacctca tgatatacctt attcccactc ctggcagcaa tatgcagaca	6240
agtctcccca tccattttgtg caacatccgg ctgcctcca cttttcagca acatcatcgc	6300
acaatcccga ctctcggctc cattcaagct tgccgcaatg tgcagtgcag tttgtcctaa	6360
aaccaatctt ccatgaaatc ttattaatct cttattaatt taatacctag tccccgccc	6420
ttcacatgaa tgtctgcacc acttcccagc agggctctcta caaccgaagc cttgccagat	6480
tgaaccgcta cgtggagagc ggtgtagtgt tctcgtgtac ggacatctac attagtagcc	6540
cgagcaatga gcattttgac gacgtcgttg aagccagcag ctgctgcgga gtgaagacct	6600
agggctcctt ttttgttggg catgaagagg gggactcctg gaagttagaa ttaacaatgt	6660
aagtcgaggg ggtgctgaga cctgtaaac ctacctctct tcaaaaacgc caatgctgtg	6720
ctagtatgct ctgaacatgc ggcaatatgc agaagcgtcg acccatcacg ggtcctagcg	6780
cgaattgagc caccaaaact gtcaattagt gactcgacca tcgaagtgtc acctcgtctc	6840
gctgcaacct gtaccggagt cttgtcctcc ttatcatgga tgttggcgtc ggcgaggagt	6900
ttgaacatga tttttagcat attttgatct ccgacttcgg ctacctggaa aattggagat	6960
agagatactg tatgtgtgca gaggcataaa ttcagatagg agtagtacca agctttgatg	7020
gagcatgaat ctagttaagg tgtatcaggg atactgtaaa ggtacggtag tccggcatat	7080
tgtatttctg acaaatctac tgtattgggt acagtaagct cagtaacctt tctgtgtacc	7140
cgttacagtg aggcaagcta aacttaggcc atttttcctg ttaaaaaacc catttaaatg	7200
ttgcctagat cagaacaagc ctcgaaatct acagcttcat cagcaaaatt tcagcttcag	7260
gagctactta aagtttcaat ttccaccctt taacctacct catgtagcgg cgtccttccc	7320
acctattctt gcacattcgc attatcacat ccagccgcaa tcgctgtccg aaccgcttcg	7380
atattcccac tccgagcggc caaatgaagc aaggtatccc cgtttccatc agctttcctg	7440
gtttgttcat ccgaaggccc acttagcaga agctccacaa tattaacatt cccaaaactg	7500
aatgccaaat gtatcggcaa ggatccatcc ccatcctctg ccattccttg atcagtatct	7560
tccaaaaatc gcttcacaat tggaaatgct ttcttggatt ttctctcgca agccacatgg	7620
attgccagct gcttttttag ccccgacct tttcggagca gctcagagta tcgcttgagg	7680
ataagctcaa gagtttcaac tccggagtac atggcggcaa tatgagtcgc gttacggcca	7740
tctttagtgc tatagttcac tcgagcacct tttcggatca tcttgtctac gatttgatcc	7800
ttgccagctt tgacgctag gaggaaggcg gtgaagccgt gctgcaagga gaatttttag	7860
aaaatggcgg gtacaatcta aagtgaaat ctaagtcagt ttcggggaat tttgggttag	7920
ggctgctaaa cggctcggag gggctcagca cattgaaaaa cgcagtgcta tatgtagtgt	7980
ttttgcagcc cgggggttcc gcaggctca cgcactagc caccatggtc ctatgtatag	8040
tgcctgctgg aaccccgaat gtgtcggcgg ctgccaaaaca tctgcctatt gactgcatt	8100
gtccaatgag aaggctcaac cccactgaag gtactacccc ctaatagtca gcagccctaa	8160
tttgggtcaa accctaaaat tgcgaacttc accgacttgt ccgagttaca gcggaaaaaa	8220
cttacattat cagccatact aaaatcactc cgcttgatag tctctatctc agactccaca	8280
ttcgccact catctctctt cgcgaaatc aaaatcctcg tctgaggatc cgcattgce	8340
aagtcctcac tcgacatttc ctcatgagac gatgcgtggg aggtgagact ctctcgaac	8400
agaggtttcc cgagaagacg atccggcggg gtgactgaat cacgggatgg ttgtttcgg	8460

-continued

```

acgaagatga tccgtgagtt ctttccgatt tggagatggg tcgaggatcg gcggaggggt 8520
ggtcgggtcag ttgggatggt gtcggtggtg aggaggtcct ggaagtggg tagaattagt 8580
tttcgtaagc ttccaggcgt gcctacacgc cttcctgttg cctacgaaaa gtccctgaatc 8640
taaaaagcat ttttggcagc atccatctaa aaaaatcggg atctttgagt agttttaaac 8700
agtgttcttc cacgaaaaaa gttttccacg tcttgccctaa gtaagcctaa gcctcagctt 8760
aagcctaagc atatgcctaa gcctaatact aagcctaagc ctgagtctga gcctgagcct 8820
aagcctattc caaagcttaa accgaagctt aagtctaggc cttagcctaa acctaagcct 8880
aaacctaagc ctaagcctaa gcctaagcct caacctaagc ctaaacctaa acctaataca 8940
atgcctacct ttttcccggt aaaccactcg gcccggtgca ccgacgtcga gcgggtttcc 9000
cgtttccgca cagttagaca tttttccgat cttgacattt tcagtattac cagaacagaa 9060
aaagaagggg aaataataca tttctctcaa ctaattgggg ggcggacgca catggtgtcc 9120
tccaacccat aaaaaagtac gaatgtgggc gattaattgc gaaaaatgcg cgaatttat 9180
ttacgactga cgacgagaag cattaactt ttggtaaagg gtgctgtggg ggtactttgg 9240
tgaaaatata gctaaaattt aggcttgggc ttgggcttag gcttaggctt aggtttcagc 9300
tcaggcttag gcttcggctc aggctttggc gtaggcttaa actttggctt aggtttaagc 9360
ttaggcttag gcttaggctt agtcttaggc ttaggcttag gcttaggctt aggctcaggt 9420
ttaagcttag acttaggctc aggtttaggc ttggcgtcag tggcgcgct tactgaagtg 9480
atatttaate actctgatga tatttaattc cgatgattaa tccacttttc tttttctcac 9540
atztatgaac caagttctaa attaaggtgg gatattttaa ggtgtgttaa catatgatat 9600
ttatttttta atttaaatat agtttctctt tttgcttctt tttataagtt ttgttaatga 9660
acgcatagtt tacaaccgcc tcgctcaaat gtattttgat aaaagtgcgc tattaggctt 9720
aagcgtcgcc ataccgccg tggtgtcata aggaattc 9758

```

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 1709

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Caenorhabditis elegans

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: amino acid sequence derived from nompC genomic sequence

&lt;400&gt; SEQUENCE: 6

```

Met Ser Arg Ser Glu Lys Cys Leu Thr Val Arg Lys Arg Glu Thr Arg
 1           5           10           15
Ser Thr Ser Val Thr Arg Ala Glu Trp Phe Thr Gly Lys Lys Met Asp
 20           25           30
Ala Ala Lys Asn Ala Phe Asp Leu Thr Thr Asp Thr Ile Pro Thr
 35           40           45
Asp Arg Pro Pro Leu Arg Arg Ser Ser Thr His Leu Gln Ile Gly Lys
 50           55           60
Asn Ser Arg Ile Ile Phe Val Pro Lys Gln Pro Ser Arg Asp Ser Val
 65           70           75           80
Thr Pro Pro Asp Arg Leu Leu Gly Lys Pro Leu Phe Arg Glu Ser Leu
 85           90           95
Thr Ser His Ala Ser Ser His Glu Glu Met Ser Ser Glu Asp Leu Ala
100          105          110
Met Ala Asp Pro Gln Thr Lys Ile Leu Tyr Phe Ala Lys Arg Asp Glu
115          120          125

```

-continued

---

Trp Ala Asn Val Glu Ser Glu Ile Glu Thr Ile Lys Arg Ser Asp Phe  
 130 135 140  
 Ser Met Ala Asp Asn His Gly Phe Thr Ala Phe Leu Leu Ala Val Lys  
 145 150 155 160  
 Ala Gly Lys Asp Gln Ile Val Asp Lys Met Ile Arg Lys Gly Ala Arg  
 165 170 175  
 Val Asp Tyr Ser Thr Lys Asp Gly Arg Asn Ala Thr His Ile Ala Ala  
 180 185 190  
 Met Tyr Ser Gly Val Glu Thr Leu Glu Leu Ile Leu Lys Arg Tyr Ser  
 195 200 205  
 Glu Leu Leu Arg Lys Gly Ala Gly Pro Lys Lys Gln Leu Ala Ile His  
 210 215 220  
 Val Ala Cys Glu Arg Lys Ser Lys Lys Ala Phe Pro Ile Val Lys Arg  
 225 230 235 240  
 Ile Leu Glu Asp Thr Asp Gln Arg Met Ala Glu Asp Gly Asp Gly Ser  
 245 250 255  
 Leu Pro Ile His Leu Ala Phe Lys Phe Gly Asn Val Asn Ile Val Glu  
 260 265 270  
 Leu Leu Leu Ser Gly Pro Ser Asp Glu Gln Thr Arg Lys Ala Asp Gly  
 275 280 285  
 Asn Gly Asp Thr Leu Leu His Leu Ala Ala Arg Ser Gly Asn Ile Glu  
 290 295 300  
 Ala Val Arg Thr Ala Ile Ala Ala Gly Cys Asp Asn Ala Asn Val Gln  
 305 310 315 320  
 Asn Arg Val Gly Arg Thr Pro Leu His Glu Cys Leu Thr Val Thr Gly  
 325 330 335  
 Thr Gln Lys Gly Tyr Val Ala Glu Val Gly Asp Gln Asn Met Leu Lys  
 340 345 350  
 Ile Met Phe Lys Leu Arg Ala Asp Ala Asn Ile His Asp Lys Glu Asp  
 355 360 365  
 Lys Thr Pro Val His Val Ala Ala Glu Arg Gly Asp Thr Ser Met Val  
 370 375 380  
 Glu Ser Leu Ile Asp Lys Phe Gly Gly Ser Ile Arg Ala Arg Thr Arg  
 385 390 395 400  
 Asp Gly Ser Thr Leu Leu His Ile Ala Ala Cys Ser Gly His Thr Ser  
 405 410 415  
 Thr Ala Leu Ala Phe Leu Lys Arg Val Pro Leu Phe Met Pro Asn Lys  
 420 425 430  
 Lys Gly Ala Leu Gly Leu His Ser Ala Ala Ala Ala Gly Phe Asn Asp  
 435 440 445  
 Val Val Lys Met Leu Ile Ala Arg Gly Thr Asn Val Asp Val Arg Thr  
 450 455 460  
 Arg Asp Asn Tyr Thr Ala Leu His Val Ala Val Gln Ser Gly Lys Ala  
 465 470 475 480  
 Ser Val Val Glu Thr Leu Leu Gly Ser Gly Ala Asp Ile His Val Lys  
 485 490 495  
 Gly Gly Glu Leu Met Asp Gly Glu Thr Cys Leu His Ile Ala Ala Arg  
 500 505 510  
 Ser Gly Asn Lys Asp Ile Met Leu Leu Leu Asp Glu Asn Ala Asp Ser  
 515 520 525  
 Lys Ile Ser Ser Lys Ile Gly Glu Thr Pro Leu Gln Val Ala Ala Lys  
 530 535 540  
 Ser Cys Asn Phe Glu Ala Ala Ser Met Ile Leu Lys His Leu Ser Glu

-continued

545	550					555					560				
Val Leu Thr	Gln Glu	Gln Leu	Lys Glu	His Val	Asn His	Arg Thr	Asn								
	565			570			575								
Asp Gly Phe	Thr Ala	Leu His	Tyr Ala	Ala Ala	Glu Ile	Glu Gln	Arg Gln								
	580			585			590								
Leu His Phe	Pro Gly	Glu Asp	Ala Lys	Leu Val	Asn Leu	Leu Leu	Ile Asp								
	595		600				605								
Tyr Gly Gly	Met Val	Glu Met	Pro Ser	Leu Asn	Ala Asn	Glu Thr	Ala								
	610			615			620								
Met His Met	Ala Ala	Arg Ser	Gly Asn	Gln Ala	Val Leu	Leu Ala	Met								
	625		630				635							640	
Val Asn Lys	Ile Gly	Ala Gly	Ala Val	Gln Ile	Val Gln	Asn Lys	Gln								
	645			650			655								
Ser Lys Asn	Gly Trp	Ser Pro	Leu Leu	Glu Ala	Cys Ala	Arg Gly	His								
	660			665			670								
Ser Gly Val	Ala Asn	Ile Leu	Leu Lys	Val Leu	Val Leu	Cys Val	Gly								
	675		680				685								
Pro Gly Pro	Gly Pro	Gly Pro	Arg Leu	Gln Gly	Arg Gly	Tyr Trp	Thr								
	690		695				700								
Arg Thr Arg	Ala Arg	Val Thr	Val Pro	Trp Leu	Gln Tyr	Gln Gly	Tyr								
	705			710			715							720	
Trp Ala Arg	Thr Arg	Thr Arg	Thr Arg	Ala Arg	Ala Thr	Gly Pro	Gly								
	725			730			735								
Leu Gln Asp	Gln Gly	Tyr Trp	Ala Arg	Thr Arg	Thr Arg	Thr Lys	Val								
	740			745			750								
Thr Val Pro	Arg Leu	Leu Leu	Gly Asp	His His	Ala Arg	Ile Asp	Val Phe								
	755		760				765								
Asp Glu Met	Gly Arg	Thr Ala	Leu His	Leu Ala	Ala Phe	Asn Gly	His								
	770		775				780								
Leu Ser Leu	Val His	Leu Leu	Leu Gln	His Lys	Ala Phe	Val Asn	Ser								
	785		790				800								
Lys Ser Lys	Thr Gly	Glu Ala	Pro Leu	His Leu	Ala Ala	Gln His	Gly								
	805			810			815								
His Val Lys	Val Val	Asn Val	Leu Val	Gln Asp	His Gly	Ala Ala	Leu								
	820			825			830								
Glu Ala Ile	Thr Leu	Asp Asn	Gln Thr	Ala Leu	His Phe	Ala Ala	Lys								
	835		840				845								
Phe Gly Gln	Leu Ala	Val Ser	Gln Thr	Leu Leu	Ala Leu	Gly Ala	Asn								
	850		855				860								
Pro Asn Ala	Arg Asp	Asp Lys	Gly Gln	Thr Pro	Leu His	Leu Ala	Ala								
	865		870				875							880	
Glu Asn Asp	Phe Pro	Asp Val	Val Lys	Leu Phe	Leu Lys	Met Arg	Asn								
	885			890			895								
Asn Asn Arg	Ser Val	Leu Thr	Ala Ile	Asp His	Asn Gly	Phe Thr	Cys								
	900			905			910								
Ala His Ile	Ala Ala	Met Lys	Gly Ser	Leu Ala	Val Val	Arg Glu	Leu								
	915		920				925								
Met Met Ile	Asp Lys	Pro Met	Val Ile	Gln Ala	Lys Thr	Lys Thr	Leu								
	930		935				940								
Glu Ala Thr	Thr Leu	His Met	Ala Ala	Ala Ala	Gly Gly	His Ala	Asn Ile								
	945		950				955							960	
Val Lys Ile	Leu Leu	Glu Asn	Gly Ala	Asn Ala	Glu Asp	Glu Asn	Ser								
	965			970			975								

-continued

---

Gly Met Thr Ala Leu His Leu Gly Ala Lys Asn Gly Phe Ile Ser Ile  
 980 985 990

Leu Glu Ala Phe Asp Lys Ile Leu Trp Lys Arg Cys Ser Arg Lys Thr  
 995 1000 1005

Gly Leu Asn Ala Leu His Ile Ala Ala Phe Tyr Gly Asn Ser Asp Phe  
 1010 1015 1020

Val Asn Glu Met Leu Lys His Val Gln Ala Thr Val Arg Ser Glu Pro  
 1025 1030 1035 1040

Pro Ile Tyr Asn His His Val Asn Lys Glu Phe Ser Thr Glu Tyr Gly  
 1045 1050 1055

Phe Thr Pro Leu His Leu Ala Ala His Ser Gly His Asp Ser Leu Val  
 1060 1065 1070

Arg Met Leu Leu Asn Gln Gly Val Gln Val Asp Ala Thr Ser Thr Thr  
 1075 1080 1085

Met Met Ser Glu Lys Glu Lys Glu Arg Ala Lys Asp Leu Leu Asn Val  
 1090 1095 1100

Ala Val Phe Ser Glu Asn Met Ala Val Glu Leu Ile Thr Ala Thr  
 1105 1110 1115 1120

Glu Tyr Asn Ala Ala Leu Leu Leu Lys Ala Lys Asp Asn Arg Gly Arg  
 1125 1130 1135

Pro Leu Leu Asp Val Leu Ile Glu Asn Glu Gln Lys Glu Val Val Ser  
 1140 1145 1150

Tyr Ala Ser Val Gln Arg Tyr Leu Thr Glu Val Trp Thr Ala Arg Val  
 1155 1160 1165

Asp Trp Ser Phe Gly Lys Phe Val Ala Phe Ser Leu Phe Val Leu Ile  
 1170 1175 1180

Cys Pro Pro Ala Trp Phe Tyr Phe Ser Leu Pro Leu Asp Ser Arg Ile  
 1185 1190 1195 1200

Gly Arg Ala Pro Ile Ile Lys Phe Val Cys His Ile Val Ser His Val  
 1205 1210 1215

Tyr Phe Thr Ile Leu Leu Thr Ile Val Val Leu Asn Ile Thr His Lys  
 1220 1225 1230

Tyr Glu Val Thr Ser Val Val Pro Asn Pro Val Glu Trp Leu Leu Leu  
 1235 1240 1245

Leu Trp Leu Ser Gly Asn Leu Val Ser Glu Leu Ser Thr Val Gly Gly  
 1250 1255 1260

Gly Ser Gly Leu Gly Ile Val Lys Val Leu Ile Leu Val Leu Ser Ala  
 1265 1270 1275 1280

Met Ala Ile Ala Val His Val Leu Ala Phe Leu Leu Pro Ala Val Phe  
 1285 1290 1295

Leu Thr His Leu Asp Asn Asp Glu Lys Leu His Phe Ala Arg Thr Met  
 1300 1305 1310

Leu Tyr Leu Lys Asn Gln Leu Phe Ala Phe Ala Leu Leu Phe Ala Phe  
 1315 1320 1325

Val Glu Tyr Leu Asp Phe Leu Thr Val His His Leu Phe Gly Pro Trp  
 1330 1335 1340

Ala Ile Ile Ile Met Tyr Asp Leu Ala Arg Phe Leu Val Ile Leu Met  
 1345 1350 1355 1360

Leu Phe Val Ala Gly Phe Thr Leu His Val Thr Ser Ile Phe Gln Pro  
 1365 1370 1375

Ala Tyr Gln Pro Val Asp Glu Asp Ser Ala Glu Leu Met Arg Leu Ala  
 1380 1385 1390

-continued

---

Ser Pro Ser Gln Thr Leu Glu Met Leu Phe Phe Ser Leu Phe Gly Leu  
1395 1400 1405

Val Glu Pro Asp Ser Met Pro Pro Leu His Leu Val Pro Asp Phe Ala  
1410 1415 1420

Lys Ile Ile Leu Lys Leu Leu Phe Gly Ile Tyr Met Met Val Thr Leu  
1425 1430 1435 1440

Ile Val Leu Ile Asn Leu Leu Ile Ala Met Met Ser Asp Thr Tyr Gln  
1445 1450 1455

Arg Ile Gln Ala Gln Ser Asp Lys Glu Trp Lys Phe Gly Arg Ala Ile  
1460 1465 1470

Leu Ile Arg Gln Met Asn Lys Lys Ser Ala Thr Pro Ser Pro Ile Asn  
1475 1480 1485

Met Leu Thr Lys Leu Ile Ile Val Leu Arg Val Ala Trp Arg Asn Arg  
1490 1495 1500

Gly Lys Ala Pro Leu Ser Thr Pro Leu Ala Ser Phe Arg Cys Met Thr  
1505 1510 1515 1520

Arg Lys Ala Gln Asp Asp Leu Arg Phe Glu Glu Asn Ile Asp Ala Phe  
1525 1530 1535

Ser Met Gly Gly Gly Gln Gln Gly Arg Gln Ser Pro Thr Asn Glu Gly  
1540 1545 1550

Arg Gly Gln Gln Glu Leu Gly Asn Ser Ala Asp Trp Asn Ile Glu Thr  
1555 1560 1565

Val Ile Asp Trp Arg Lys Ile Val Ser Met Tyr Tyr Gln Ala Asn Gly  
1570 1575 1580

Lys Leu Thr Asp Gly Arg Thr Lys Glu Asp Val Asp Leu Ala Met Ala  
1585 1590 1595 1600

Val Pro Thr Ser Phe Ile Lys Pro Gln Gly Pro Asp Thr Thr Cys Arg  
1605 1610 1615

Pro Ile Asp Tyr Thr Trp Leu Arg Leu Cys Lys Thr Lys Ser His Gly  
1620 1625 1630

Ser Gly Leu Ser Ile Val Arg Arg Lys Thr Arg Gly Lys Ile Val Tyr  
1635 1640 1645

Ser Thr Arg Thr Asn Thr Ser Val Leu Gln Ile Asn Ser Ser Arg Asn  
1650 1655 1660

Ala Pro Lys Ile Tyr Leu Arg Tyr Gly Arg Ala Lys Ile Ala His Phe  
1665 1670 1675 1680

Phe Phe Thr Ser Thr Thr Leu Lys Gly Gly Ala Phe Met Trp His Gly  
1685 1690 1695

Leu Ala Ala Arg Leu Cys Lys Ile Arg Val Asp His Met  
1700 1705

<210> SEQ ID NO 7  
<211> LENGTH: 12  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: amino acid  
sequence conserved between Drosophila and C.  
elegans encoding degenerate primer sets

<400> SEQUENCE: 7

Leu Asp Val Leu Ile Glu Asn Glu Gln Lys Glu Val  
1 5 10

<210> SEQ ID NO 8  
<211> LENGTH: 11  
<212> TYPE: PRT

-continued

---

<213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:amino acid  
 sequence conserved between Drosophila and C.  
 elegans encoding degenerate primer sets

<400> SEQUENCE: 8

His His Leu Phe Gly Pro Trp Ala Ile Ile Ile  
 1 5 10

<210> SEQ ID NO 9  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:amino acid  
 sequence conserved between Drosophila and C.  
 elegans encoding degenerate primer sets

<400> SEQUENCE: 9

Val Leu Ile Asn Leu Leu Ile Ala Met Met Ser Asp Thr Tyr Gln Arg  
 1 5 10 15

Ile Gln

<210> SEQ ID NO 10  
 <211> LENGTH: 19  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #1

<400> SEQUENCE: 10

Ile Leu Leu Leu Leu Val Ala Phe Ile Val Cys Pro Pro Val Trp Ile  
 1 5 10 15

Gly Phe Thr

<210> SEQ ID NO 11  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #2

<400> SEQUENCE: 11

Tyr Trp Tyr Glu Val Gly Leu Leu Ile Trp Leu Ser Gly Leu Leu Leu  
 1 5 10 15

Phe Glu Leu Thr  
 20

<210> SEQ ID NO 12  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #3

<400> SEQUENCE: 12

Ile Lys Val Leu Val Leu Leu Leu Gly Met Ala Gly Val Gly Val His  
 1 5 10 15

Val Ser Ala Phe  
 20

-continued

---

<210> SEQ ID NO 13  
 <211> LENGTH: 25  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #4

&lt;400&gt; SEQUENCE: 13

Thr Leu Val Tyr Cys Arg Asn Gln Cys Phe Ala Leu Ala Phe Leu Leu  
 1 5 10 15  
 Ala Cys Val Gln Ile Leu Asp Phe Leu  
 20 25

<210> SEQ ID NO 14  
 <211> LENGTH: 20  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #5

&lt;400&gt; SEQUENCE: 14

Phe Leu Ala Val Leu Ala Ile Phe Val Phe Gly Phe Ser Met His Ile  
 1 5 10 15  
 Val Ala Leu Asn  
 20

<210> SEQ ID NO 15  
 <211> LENGTH: 23  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #6

&lt;400&gt; SEQUENCE: 15

Ile Val Phe Gly Ile Tyr Met Leu Val Ser Val Val Val Leu Ile Asn  
 1 5 10 15  
 Leu Leu Ile Ala Met Met Ser  
 20

<210> SEQ ID NO 16  
 <211> LENGTH: 17  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #7

&lt;400&gt; SEQUENCE: 16

Tyr Ile Asn Phe Ile Leu His Cys Val Leu Ile Ile Leu Tyr Phe Ser  
 1 5 10 15  
 Ile

<210> SEQ ID NO 17  
 <211> LENGTH: 19  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence:nompC  
 transmembrane domain (channel region) #8

&lt;400&gt; SEQUENCE: 17



-continued

---

Ile Tyr Leu Met Ile His Leu Ser Ile Val Gly Ile Thr Pro Ile Tyr  
 1                    5                    10                    15

Pro Val Leu

---

What is claimed is:

1. An in vitro method for identifying a compound that modulates mechanosensory receptor activity in eukaryotic cells, the method comprising the steps of:

- (i) contacting the compound with a mechanosensory receptor protein, the protein having cation channel activity and comprising a sequence with 90% or greater identity to the sequence of SEQ ID NO:4; and
- (ii) determining the functional effect of the compound on the mechanosensory receptor protein, thereby identifying a compound that modulates said mechanosensory receptor activity.

2. The method of claim 1, wherein the mechano sensory receptor protein is expressed in a eukaryotic cell or cell membrane.

10 3. The method of claim 2, wherein the functional effect is determined by detecting a change in the membrane potential of the cell or cell membrane.

15 4. The method of claim 2, wherein the functional effect is determined by detecting a change in an intracellular ion concentration.

5. The method of claim 4, wherein the ion is selected from the group consisting of K<sup>+</sup> and Ca<sup>2+</sup>.

20 6. The method of claim 1, wherein the protein comprises the amino acid sequence of SEQ ID NO:4.

7. The method of claim 1, wherein the protein is recombinant.

\* \* \* \* \*